

INFORMATION
ANALYSIS
DIRECTORATE



Health Inequalities

Annual Report 2024

A product of the NI Health and Social Care Inequalities Monitoring System

Annual Publication 2024



Department of
Health

An Roinn Sláinte

Mánnystrie O Poustie

www.health-ni.gov.uk



NISRA

Northern Ireland
Statistics and Research Agency
Gníomhaireacht Thuaisceart Éireann
um Staitisticí agus Taighde

Annual Publication: March 2024

Health Inequalities

Annual Report 2024

Authors: Robbyn Atcheson & Caolan Laverty

Public Health Information & Research Branch, Information Analysis Directorate

Lead Statistician: Bill Stewart

Department of Health
Castle Buildings, Belfast BT4 3SQ

☎ Tel: 02890 522 501

☎ Ext: 22501

✉ Email: healthinequalities@health-ni.gov.uk

<http://www.health-ni.gov.uk/topics/dhssps-statistics-and-research/health-inequalities-statistics>

For information on other Government statistics contact:
The Northern Ireland Statistics and Research Agency (NISRA)
Colby House, Stranmillis Court, Belfast, BT9 5RR
☎ Tel: 02890388400 ✉ Email: info@nisra.gov.uk
www.nisra.gov.uk

© Crown Copyright 2024. Published with the permission of the Controller of Her Majesty's Stationary Office.

This publication is Crown copyright and may be reproduced free of charge in any format or medium.
Any material used must be acknowledged and sourced to the NI Health and Social Care Inequalities Monitoring System.

Information Analysis Directorate (IAD) sits within the **Department of Health (DoH)** and carries out various statistical work and research on behalf of the department. It comprises four statistical areas: Hospital Information, Community Information, Public Health Information & Research and Project Support Analysis.

IAD is responsible for compiling, processing, analysing, interpreting and disseminating a wide range of statistics covering health and social care.

The statisticians within IAD are out-posted from the Northern Ireland Statistics & Research Agency (NISRA) and our statistics are produced in accordance with the principles and protocols set out in the UK Code of Practice for Official Statistics.

About Public Health Information and Research Branch

The role of Public Health Information and Research Branch (PHIRB) is to support public health policy development through managing the public health survey function while also providing analysis and monitoring data. The head of the branch is the Principal Statistician, Mr. Bill Stewart.

In support of the public health survey function, PHIRB is involved in the commissioning, managing and publishing of results from departmental funded surveys, such as the Health Survey Northern Ireland, All Ireland Drug Prevalence Survey, Young Persons Behaviour & Attitudes Survey, Patient Experience Surveys and the Adult Drinking Patterns Survey.

The branch also houses the NI Health and Social Care Inequalities Monitoring System which covers a range of different health inequality/equality based projects conducted for both the region as well as for more localised area levels. In addition, PHIRB is responsible for the production of official life expectancy estimates for NI, and areas within the region.

PHIRB provides support to a range of key DoH NI strategies including Making Life Better, a 10 year cross-departmental public health strategic framework as well as a range of other departmental strategies such as those dealing with suicide, sexual health, breastfeeding, tobacco control and obesity prevention. It also has a key role in supporting the Substance Use Strategy, by maintaining and developing key departmental databases such as, the Substance Misuse Database, Impact Measurement Tool and the Census of Drug & Alcohol Treatment Services, which are all used to monitor drug misuse and treatments across Northern Ireland. In addition to Departmental functions, PHIRB also support the executive level Programme for Government and its strategic outcomes through a series of performance indicators.

CONTENTS

	PAGE NUMBER
<u>KEY FINDINGS</u>	5
INTRODUCTION	7
FORMAT OF THE REPORT	7
REGIONAL HEALTH INEQUALITIES	12
SUMMARY OF CHANGES IN REGIONAL INEQUALITY GAPS	13
LIFE EXPECTANCY & GENERAL HEALTH	15
PREMATURE MORTALITY	18
MAJOR DISEASES	21
HOSPITAL ACTIVITY	24
MENTAL HEALTH	26
ALCOHOL, SMOKING & DRUGS	28
PREGNANCY & EARLY YEARS	31
DIET & DENTAL HEALTH	33
SUB-REGIONAL HEALTH INEQUALITIES	37
<u>HEALTH & SOCIAL CARE TRUSTS</u>	38
COMPARISON OF TRUST HEALTH OUTCOMES AGAINST REGIONAL AVERAGE	38
LARGEST DEPRIVATION INEQUALITY GAPS IN EACH TRUST	38
BELFAST	39
NORTHERN	41
SOUTH EASTERN	43
SOUTHERN	45
WESTERN	47
<u>LOCAL GOVERNMENT DISTRICTS (LGD)</u>	49
COMPARISON OF LGD HEALTH OUTCOMES AGAINST REGIONAL AVERAGE	49
LARGEST DEPRIVATION INEQUALITY GAPS IN EACH LGD	50
ANTRIM & NEWTOWNABBIEY	51
ARDS & NORTH DOWN	53
ARMAGH CITY, BANBRIDGE & CRAIGAVON	55
BELFAST LGD	57
CAUSEWAY COAST & GLENS	59
DERRY CITY & STRBANE	61
FERMANAGH & OMAGH	63
LISBURN & CASTLEREAGH	65
MID & EAST ANTRIM	67
MID ULSTER	69
NEWRY, MOURNE & DOWN	71
APPENDICIES	
APPENDIX A: SOCIAL GRADIENT OF HEALTH	73
APPENDIX B: POPULATION ATTRIBUTABLE RISK (PAR) OF DEPRIVATION	83
APPENDIX C: ADDITIONAL INDICATORS	84
APPENDIX D: URBAN RURAL ANALYSIS	87
APPENDIX E: TECHNICAL NOTES & DEFINITIONS	88

KEY FINDINGS – REGIONAL (NI)

- Deprivation gaps for male and female life expectancies saw no change since 2016-18 with the most-least deprivation gaps stood at 7.2 years for males and 4.8 years for females in 2020-22. While female life expectancy at birth remained similar in all areas over the last five years, male life expectancy decreased in 2020-22 in NI and the most and least deprived areas.
- Inequality gaps narrowed for both male and female disability-free life expectancies (DFLE) following improvements in the most deprived areas. There were no changes in inequality gaps observed for male or female healthy life expectancies (HLE) since 2016-18, with the most-least deprived gaps in HLE stood at 12.2 years for males and 14.2 years for females in 2020-22.
- Large inequality gaps continue to highlight markedly higher rates of premature mortality in the most deprived areas, with none of the gaps showing a notable change over the analysed period, with the exception of preventable mortality where the gap widened as the rate in the most deprived areas increased to now treble the rate in the least deprived areas.
- Large inequality gaps continue to exist for mental health indicators. Prescription rates for mood and anxiety disorders increased regionally and for most & least deprived areas between 2018 and 2022, with the rate in the most deprived areas 66% higher than in the least deprived areas. In 2020-22 the suicide mortality rate in the most deprived areas was more than two and a half times the rate observed in the least deprived areas.
- Alcohol and drug related indicators continue to show some of the largest health inequalities monitored in NI. The deprivation gap for drug misuse deaths widened over the analysed period and showed the largest inequality gap, where mortality in the most deprived areas was almost six times that of the least deprived.
- While improvements were observed regionally and in the most and least deprived areas with the under 20 teenage birth rate, in addition to regionally and in the most deprived areas for the proportion of mothers reporting smoking, the relative inequality gaps remain very large; with rates in the most deprived areas being over five times that in the least deprived in 2022.
- In 2022/23 the percentage of primary 1 pupils in the most deprived areas affected by obesity was more than double the proportion in the least deprived areas. The inequality gap in year 8 pupils affected by obesity was slightly lower, with the proportion in the most deprived areas 94% higher than in the least deprived areas.

KEY FINDINGS – SUB-REGIONAL (HSC TRUST & LGD)

- Male and female life expectancies were lowest in the Belfast LGD (75.8 years and 80.4 years respectively). Male life expectancy was highest in Lisburn & Castlereagh LGD (80.4 years), and female life expectancy was highest in Lisburn & Castlereagh and Mid Ulster LGDs (83.1 years).
- Similar to the regional picture, deprivation related inequality was most prominent in indicators relating to alcohol, drugs, self-harm, smoking during pregnancy and teenage births, which were among the five largest inequality gaps for the majority of Trusts and LGDs.
- Drug misuse mortality was the largest inequality gap in three of the five HSC Trusts. In the Western Trust, the rate in its most deprived areas was almost two and a half times (145%) the Trust average.
- Drug related deaths was the largest inequality gap in five of the 11 LGDs, and in the Mid & East Antrim LGD the rate of drug related mortality in the most deprived areas was more than treble (220%) the LGD average.
- Large inequality gaps for alcohol related admissions also exist in the majority of Trusts and LGDs. The rate in their most deprived areas was at least double the Trust/LGD average for the Western Trust (115%), Belfast Trust (100%), Derry City & Strabane LGD (116%), and Mid & East Antrim LGD (120%).
- Alcohol specific mortality showed the largest gap in the Causeway Coast and Glens LGD (111%), and Newry, Mourne & Down LGD (68%) whilst alcohol related admissions showed the largest gap in the Mid Ulster LGD (59%).
- The under 20 teenage birth rate was the largest inequality gap in two LGDs: Armagh City, Banbridge & Craigavon (134%) and Lisburn & Castlereagh (155%).
- Drug related admissions was the largest inequality gap in the Ards & North Down (97%) and self-harm admissions was the largest inequality gap observed in the Fermanagh & Omagh LGD (67%).

INTRODUCTION

This annual publication is one of a series of reports produced as part of the NI Health & Social Care Inequalities Monitoring System (HSCIMS) and presents a comprehensive analysis of health outcomes and inequality gaps between the most and least deprived areas of NI, and within Health & Social Care (HSC) Trust and Local Government District (LGD) areas across a range of indicators. This report is an update of the Health Inequalities Annual Report 2023. The report is accompanied by downloadable data tables¹, which contain all figures, including urban and rural breakdowns. The most recent figures reported in this release typically include data from 2021 and 2022, and therefore reflect to an extent, the impact of the coronavirus (COVID-19) pandemic. In particular, figures related to hospital admissions, dental indicators and childhood obesity have been significantly impacted due to service reconfiguration and/or restrictions.

ACCOUNTABILITY STATEMENT

Our statistical practice is regulated by the Office for Statistics Regulation (OSR). OSR sets the standards of trustworthiness, quality and value in the Code of Practice for Statistics that all producers of official statistics should adhere to. You are welcome to contact us directly at healthinequalities@health-ni.gov.uk with any comments about how we meet these standards. Alternatively, you can contact OSR by emailing regulation@statistics.gov.uk or via the OSR website.

OFFICIAL STATISTICS

This report is an official statistics publication. Official statistics are statistics produced by Crown bodies and other organisations listed within an Official Statistics Order, on behalf of the UK government or devolved administrations. They provide a factual basis for assessment and decisions on economic, social and environmental issues at all levels of society. This broad definition of official statistics means that the scope of official statistics can be adapted over time to suit changing circumstances.

FORMAT OF THE REPORT

This report is separated into two sections, the first focusing on regional health inequalities and the second presenting sub-regional analysis. The regional section contains separate chapters for each theme/topic area, with each section containing a summary of the key findings, followed by individual indicator analysis. The sub-regional section presents a condensed summary of findings for each HSC Trust and LGD accompanied by downloadable data tables¹ which contain all figures.

ASSESSMENT OF CHANGE OVER TIME

An indication of the changes observed at the NI level has been provided for each health outcome in addition to assessments of the changes in the rates in the most deprived and least deprived areas, and in the most-least deprived inequality gap. An improvement or decline in the rate is only indicated when the change is statistically significant, or where there is a clear and consistent trend observed over the series. For a notable change in the inequality gap to have occurred, a significant change in at least one of the areas (most/least deprived) has to have been observed, or, where no statistically significant change is apparent then a change in the gap will have been deemed to have

¹ <https://www.health-ni.gov.uk/articles/health-inequalities-statistics>

occurred if there is a clear and consistent trend in both the outcome/activity and the gap over the analysed period².

It should be noted that inequality gaps for indicators can exist in either direction; however health outcomes generally tend to be worse in the most deprived areas than in the least deprived. For the purposes of this report, a positive value for the gap means that the health outcomes experienced in the most deprived areas were worse than in the least deprived³.

² Indicated changes are based on a subjective assessment of the available data.

³ Assessments of change for outcomes relating to service-based indicators, including all hospital admission and dental treatment activity, have been analysed and presented based on whether there was an observed increase or decrease in activity, rather than positive or negative changes. See 'Service-based indicators' note on Page 8.

NOTES FOR USER

- **Regional Inequality Gaps** refer to the difference in health outcomes between the 20% most deprived and 20% least deprived areas of Northern Ireland.
- **Sub-regional Inequality Gaps** refer to the difference between health outcomes for
 - The 20% most deprived areas of an area (LGD/Trust) and the area's average
 - The Trust/LGD and the regional average.
- **Deprivation Measure:** the 20% most and least deprived areas are defined according to the Northern Ireland Multiple Deprivation Measure (NIMDM) 2017.⁴
- **Rounded Figures:** some individual figures have been rounded to either zero or one decimal place independently. As a result, the sum of component items may not therefore always add to the totals shown.
- **Additional Indicators:** figures relating to ten additional indicators such as Median Fire Response Times and Median Ambulance Response Times, which form part of the HSCIMS but are not contained in the main body of the report, can be found in [Appendix C](#). Previous reports did not present year 8 BMI indicators due to data quality limitations. This report now includes these indicators again, with single year rates presented for primary 1 and year 8 BMI.
- **Service-based indicators:** assessments of change for outcomes relating to service-based indicators, including all hospital admission and dental treatment activity, have been analysed and presented based on whether there was an observed increase or decrease in activity, rather than positive or negative changes to health outcomes. This is due to difficulties in ascertaining whether any changes in rates are due to changes in demand (i.e., health of the population), or, as a result of changes in service provision. All figures relating to these indicators should be treated with caution as they may also be impacted by external factors that are not reflective of service demand. As unmet demand is not accounted for in the data, these indicators should therefore be viewed as indicators of service provision rather than demand. In addition, sub-regional comparisons have been analysed and presented based on whether activity was higher or lower, rather than better or worse as it is not possible to ascertain whether a sub-regional gap in utilisation of services fully constitutes a difference in health outcomes due to potentially different service provision profiles across different geographic areas.
- **Further Analysis:** The appendix section included at the back of the report provides further analysis regarding the Social Gradient of Health ([Appendix A](#)) and the Population Attributable Risk (PAR) of Deprivation ([Appendix B](#)).
- **Urban/Rural Analysis:** In addition urban and rural figures for each indicator have been included within the accompanying downloadable tables, and a summary assessment of Urban-Rural gaps has been provided in [Appendix D](#).
- **District Electoral Areas (DEAs):** figures are included within the accompanying downloadable tables.

⁴ <https://www.nisra.gov.uk/statistics/deprivation/northern-ireland-multiple-deprivation-measure-2017-nimdm2017>

- **Data limitations** mean that not all 58 health indicators analysed at a regional level can be analysed at Trust, LGD or DEA level. In this report, 53 health indicators have been presented at Trust and 51 at LGD level, with 38 reported at DEA level. A full list can be found in [Table 5](#).
- For **further information** regarding the methodologies, indicator descriptions and sources of data used to produce the analyses throughout this report, please refer to [Appendix E: Technical Notes & Definitions](#).

Rebased Mid-Year Population Estimates (2011-2021) for Northern Ireland:

Figures contained within this report have been calculated using the rebased population figures released by NISRA Census Office on 29th June 2023. This series replaces the old rolled-forward series which was based on the 2011 Census. The Rebased Mid-Year Population Estimates series reflects the results of Census 2021 and revises all previous population estimates from 2011 to 2021. As a result, some figures within this report may differ from those previously published. For further information see <https://www.nisra.gov.uk/publications/2011-21-rebased-mid-year-population-estimates-northern-ireland>.

Understanding Gaps

Regional Level:

A positive inequality gap means that the health outcomes in the most deprived areas are worse than in the least deprived areas. For service-based indicators, a positive inequality gap means that activity in the most deprived areas is higher than in the least deprived areas.

Sub-regional Level:

A positive inequality gap between the Trust or LGD and its most deprived areas means that the health outcomes in the most deprived areas are worse than the Trust or LGD average. For service-based indicators, a positive inequality gap between the Trust or LGD and its most deprived areas means that activity in the most deprived areas is higher than the Trust or LGD average.

Similarly, a positive inequality gap between the Trust or LGD and NI means that the health outcomes in the Trust or LGD are worse than the NI average. For service-based indicators, a positive inequality gap between the Trust or LGD and NI means that the level of activity in the Trust or LGD is higher than the NI average.

Other routine reports in the HSCIMS series include:

Life Expectancy in Northern Ireland – presenting the latest official estimates of life expectancy in Northern Ireland and an examination of the causes that contribute to the change in life expectancy over time, as well as the differentials between genders and across Local Government Districts. The latest figures for life expectancy at 65, healthy life expectancy and disability-free life expectancy are also included. <https://www.health-ni.gov.uk/articles/life-expectancy-northern-ireland>

Making life better: monitoring the wider social determinants of health & wellbeing - key indicators – monitoring report for the key indicators of the wider social determinants of health & wellbeing, contained in the Making Life Better,⁵ the public health strategic framework for NI. <https://www.health-ni.gov.uk/articles/social-determinants-health-statistics>.

⁵ www.health-ni.gov.uk/topics/public-health-policy-and-advice/making-life-better-whole-system-strategic-framework-public

Regional Health Inequalities

Regional health inequalities refer to the difference in health outcomes between the 20% most deprived and 20% least deprived areas of Northern Ireland according to the Northern Ireland Multiple Deprivation Measure. This section contains separate chapters for each theme/topic area, with each chapter containing a summary of the key findings, followed by individual indicator analysis.

For ease of understanding, each theme is assigned a separate colour (for example blue is used for 'Life Expectancy and General Health'). Each indicator is broken down by Northern Ireland average, the average in the 20% Most Deprived areas, the 20% Least Deprived areas and the Deprivation Gap.

SUMMARY OF CHANGES IN REGIONAL INEQUALITY GAPS OVER THE LAST 5 YEARS⁶

Most-Least Deprived Inequality Gaps that Widened over the Analysed Period

6 indicators had inequality gaps that **widened** over the analysed period.

Category	Change in Health Outcomes		
	Northern Ireland	Most Deprived Areas	Least Deprived Areas
Female Life Expectancy at age 65	No Change	No Change	No Change
Standardised Death Rate - Preventable	Negative Change	Negative Change	No Change
Standardised Death Rate - Smoking Attributable	Positive Change	Positive Change	Positive Change
Standardised Death Rate - Drug Misuse	Negative Change	Negative Change	No Change
Infant Mortality Rate	No Change	Negative Change	No Change
Low Birth Weight	No Change	No Change	No Change

Most-Least Deprived Inequality Gaps that Narrowed over the Analysed Period

11 indicators had inequality gaps that **narrowed** over the analysed period.

Category	Change in Health Outcomes		
	Northern Ireland	Most Deprived Areas	Least Deprived Areas
Male Life Expectancy at Age 65	No Change	No Change	No Change
Male Disability Free Life Expectancy	Positive Change	Positive Change	No Change
Female Disability Free Life Expectancy	No Change	Positive Change	No Change
Standardised Admission Rate – Emergency Admissions	Decrease	Decrease	Decrease
Standardised Admission Rate - Self-Harm	Decrease	Decrease	Decrease
Standardised Admission Rate - Alcohol Related Causes	Decrease	Decrease	Decrease
Standardised Death Rate – Lung Cancer	Positive Change	Positive Change	No Change
Standardised Admission Rate - Drug Related Causes	Decrease	Decrease	Decrease
Breastfeeding on Discharge	Positive Change	Positive Change	No Change
Standardised Dental Extraction Rate (U18)	Decrease	Decrease	Decrease
Standardised Dental Registration Rate	Increase	Increase	Increase

⁶ There are six indicators; four relating to Primary 1 and Year 8 BMI, in addition to Small for Gestational Age and SDR – COVID-19, for which regional assessments of change in the gap has not been carried out. This is due to an absence of data for previous years, meaning a full time series is not available.

Most-Least Deprived Inequality Gaps that Showed No Notable Change over the Analysed Period

35 indicators had inequality gaps that **showed no notable change** over the period analysed

Category	Change in Health Outcomes		
	Northern Ireland	Most Deprived Areas	Least Deprived Areas
Male Life Expectancy at Birth	Negative Change	Negative Change	Negative Change
Female Life Expectancy at Birth	No Change	No Change	No Change
Male Healthy Life Expectancy	Positive Change	Positive Change	Positive Change
Female Healthy Life Expectancy	Positive Change	Positive Change	No Change
Standardised Death Rate – All Deaths	No Change	No Change	No Change
Potential Years of Life Lost	No Change	Negative Change	No Change
Standardised Death Rate - Treatable	Positive Change	Positive Change	Positive Change
Standardised Death Rate - Avoidable	No Change	No Change	No Change
Standardised Death Rate - Circulatory U75	Positive Change	Positive Change	No Change
Standardised Death Rate - Respiratory U75	Positive Change	Positive Change	No Change
Standardised Death Rate - Cancer U75	Positive Change	Positive Change	Positive Change
Standardised Death Rate - All Cause U75	No Change	No Change	No Change
Standardised Admission Rate - Circulatory	Decrease	Decrease	Decrease
Standardised Admission Rate - Circulatory U75	Decrease	Decrease	Decrease
Standardised Prescription Rate - Antihypertensive	Negative Change	Negative Change	Negative Change
Standardised Prescription Rate - Statin	Negative Change	Negative Change	Negative Change
Standardised Admission Rate - Respiratory	Decrease	Decrease	Decrease
Standardised Admission Rate - Respiratory U75	Decrease	Decrease	Decrease
Standardised Incidence Rate - Cancer	No Change	No Change	No Change
Standardised Admission Rate - All Admissions	Decrease	Decrease	Decrease
Standardised Attendance Rate - Emergency Care	Decrease	Decrease	Decrease
Standardised Admission Rate - Elective Inpatient Admissions	Decrease	Decrease	Decrease
Standardised Admission Rate - Day Case Admissions	Decrease	Decrease	Decrease
Crude Death Rate - Suicide	No Change	No Change	No Change
Standardised Prescription Rate - Mood & Anxiety	Negative Change	Negative Change	Negative Change
Standardised Death Rate - Alcohol Specific	Negative Change	Negative Change	Negative Change
Standardised Incidence Rate – Lung Cancer	No Change	No Change	No Change
Standardised Death Rate – Drug Related Causes	Negative Change	Negative Change	Negative Change
Smoking During Pregnancy	Positive Change	Positive Change	No Change
Teenage Birth Rate U20	Positive Change	Positive Change	Positive Change
Standardised Dental Filling Rate	Decrease	Decrease	Decrease
Standardised Dental Filling Rate (U18)	Decrease	Decrease	Decrease
Standardised Dental Extraction Rate	Increase	Increase	Increase
Standardised Dental Crowning Rate	Decrease	Decrease	Decrease
Standardised Dental Registration Rate (U18)	No Change	No Change	No Change

Life Expectancy and General Health

There were no changes over the analysed period in five of the nine inequality gaps examined. While the inequality gap in female life expectancy at age 65 widened, the most-least deprived gaps for male life expectancy at age 65, and for both male and female disability-free life expectancy narrowed. While female life expectancy at birth remained similar in all areas over the last five years, male life expectancy decreased in NI and the most and least deprived areas. Healthy Life Expectancy (HLE) improved for males and females in NI and also the most deprived areas, in addition to the least deprived areas for males. The standardised death rate for all deaths remained similar over the period for all areas.

Male Life Expectancy at Birth NI: Negative Change, Most Deprived: Negative Change, Least Deprived: Negative Change, Deprivation Gap: No Change

Years	Northern Ireland	Most Deprived	Least Deprived	Deprivation Gap
2016-18	78.7	74.7	81.8	7.1
2017-19	78.8	74.7	81.7	7.0
2018-20	78.7	74.5	81.4	6.9
2019-21	78.5	73.9	81.3	7.3
2020-22	78.4	74.0	81.2	7.2

Female Life Expectancy at Birth NI: No Change, Most Deprived: No Change, Least Deprived: No Change, Deprivation Gap: No Change

Years	Northern Ireland	Most Deprived	Least Deprived	Deprivation Gap
2016-18	82.4	79.7	84.1	4.4
2017-19	82.6	79.6	84.5	4.9
2018-20	82.4	79.3	84.3	5.0
2019-21	82.3	79.2	84.2	5.1
2020-22	82.3	79.3	84.1	4.8

Male Life Expectancy at Age 65 NI: No Change, Least Deprived: No Change, Most Deprived: No Change, Deprivation Gap: Narrowed

Years	Northern Ireland	Most Deprived	Least Deprived	Deprivation Gap
2016-18	18.4	16.5	19.8	3.2
2017-19	18.5	16.7	19.8	3.1
2018-20	18.5	16.6	19.7	3.1
2019-21	18.4	16.5	19.5	3.0
2020-22	18.3	16.5	19.4	2.9

Female Life Expectancy at Age 65

NI: No Change, Most Deprived: No Change, Least Deprived: No Change, Deprivation Gap: Widened

Years	Northern Ireland	Most Deprived	Least Deprived	Deprivation Gap
2016-18	20.7	19.2	21.6	2.4
2017-19	20.9	19.3	21.8	2.6
2018-20	20.7	19.0	21.7	2.7
2019-21	20.7	19.0	21.7	2.7
2020-22	20.6	18.9	21.6	2.8

Male Healthy Life Expectancy⁷

NI: Positive Change, Most Deprived: Positive Change, Least Deprived: Positive Change, Deprivation Gap: No Change

Years	Northern Ireland	Most Deprived	Least Deprived	Deprivation Gap
2016-18	59.7	51.4	65.4	14.0
2017-19	59.2	51.6	65.1	13.5
2018-20	60.1	53.8	65.8	12.0
2019-21	60.6	55.1	66.3	11.2
2020-22	61.2	54.8	67.0	12.2

Female Healthy Life Expectancy⁷

NI: Positive Change, Most Deprived: Positive Change, Least Deprived: No Change, Deprivation Gap: No Change

Years	Northern Ireland	Most Deprived	Least Deprived	Deprivation Gap
2016-18	60.8	50.8	66.0	15.2
2017-19	61.0	51.7	67.1	15.4
2018-20	62.1	52.5	67.2	14.7
2019-21	62.7	52.3	67.4	15.1
2020-22	62.7	52.9	67.1	14.2

Male Disability Free Life Expectancy⁷

NI: Positive Change, Most Deprived: Positive Change, Least Deprived: No Change, Deprivation Gap: Narrowed

Years	Northern Ireland	Most Deprived	Least Deprived	Deprivation Gap
2016-18	57.3	48.3	62.8	14.5
2017-19	57.9	49.6	62.1	12.5
2018-20	58.2	50.6	62.0	11.4
2019-21	58.6	52.4	62.2	9.8
2020-22	59.2	52.4	63.5	11.1

⁷ Since 2020/21, the Health Survey NI, from which HLE and DFLE estimates are produced, has been telephone based using a smaller sample size and has not included children. To ensure the figures remain as representative as possible of the entire population, data for children has been held constant from 2019. This should be considered when assessing changes. Please see [Appendix E](#) for more detail.

Female Disability Free Life Expectancy⁸

NI: No Change, Most Deprived: Positive Change,
Least Deprived: No Change, Deprivation Gap:
Narrowed

Years	Northern Ireland	Most Deprived	Least Deprived	Deprivation Gap
2016-18	57.2	47.8	61.7	13.9
2017-19	58.4	49.6	62.9	13.3
2018-20	59.0	50.2	62.3	12.1
2019-21	58.7	50.4	61.7	11.3
2020-22	58.5	50.9	61.9	11.0

Standardised Death Rate – All Deaths

NI: No Change, Most Deprived: No Change, Least
Deprived: No Change, Deprivation Gap: No Change

Years	Northern Ireland	Most Deprived	Least Deprived	Deprivation Gap
2018	1,009	1,272	852	49%
2019	972	1,216	839	45%
2020	1,070	1,327	946	40%
2021	1,047	1,324	871	52%
2022	1,008	1,234	866	42%

⁸ Since 2020/21, the Health Survey NI, from which HLE and DFLE estimates are produced, has been telephone based using a smaller sample size and has not included children. To ensure the figures remain as representative as possible of the entire population, data for children has been held constant from 2018/20. This should be considered when assessing changes. Please see [Appendix E](#) for more detail.

Premature Mortality

Large inequality gaps continue to highlight markedly higher rates of premature mortality in the most deprived areas. Of the inequality gaps analysed, none showed a notable change over the period, with the exception of preventable mortality where the gap widened as the rate in the most deprived areas increased to now treble the rate in the least deprived areas. For treatable mortality, rates for NI and the most and least deprived areas decreased with the most-least deprived gap remaining similar over the period. The largest inequality gap was seen for the under 75 respiratory death rate where the rate in the most deprived areas was almost three and a half times the rate in the least deprived areas.

Potential Years of Life Lost NI: No Change, Most Deprived: Negative Change, Least Deprived: No change, Deprivation Gap: No Change

Years	Northern Ireland	Most Deprived	Least Deprived	Deprivation Gap
2016-18	8.5	12.9	5.7	126%
2017-19	8.4	13.0	5.7	129%
2018-20	8.5	13.3	5.9	125%
2019-21	8.7	13.8	5.9	133%
2020-22	8.7	13.6	5.9	130%

Standardised Death Rate – Treatable NI: Positive Change, Most Deprived: Positive Change, Least Deprived: Positive Change, Deprivation Gap: No Change

Years	Northern Ireland	Most Deprived	Least Deprived	Deprivation Gap
2014-18	85	122	64	89%
2015-19	83	119	62	91%
2016-20	82	116	62	86%
2017-21	80	114	60	89%
2018-22	80	110	60	83%

Standardised Death Rate – Preventable NI: Negative Change, Most Deprived: Negative Change, Least Deprived: No Change, Deprivation Gap: Widened

Years	Northern Ireland	Most Deprived	Least Deprived	Deprivation Gap
2014-18	168	290	103	182%
2015-19	169	295	100	194%
2016-20	173	299	103	191%
2017-21	178	310	103	200%
2018-22	179	312	103	202%

Standardised Death Rate – Avoidable

NI: No Change, Most Deprived: No Change, Least Deprived: No Change, Deprivation Gap: No Change

Years	Northern Ireland	Most Deprived	Least Deprived	Deprivation Gap
2014-18	253	412	167	146%
2015-19	252	414	163	155%
2016-20	254	415	165	152%
2017-21	259	424	164	159%
2018-22	259	422	163	158%

Standardised Death Rate – Circulatory U75

NI: Positive Change, Most Deprived: Positive Change, Least Deprived: No Change, Deprivation Gap: No Change

Years	Northern Ireland	Most Deprived	Least Deprived	Deprivation Gap
2014-18	71	110	47	134%
2015-19	70	108	45	142%
2016-20	70	106	45	136%
2017-21	69	105	44	137%
2018-22	68	103	46	127%

Standardised Death Rate – Respiratory U75

NI: Positive Change, Most Deprived: Positive Change, Least Deprived: No Change, Deprivation Gap: No Change

Years	Northern Ireland	Most Deprived	Least Deprived	Deprivation Gap
2014-18	35	66	19	256%
2015-19	35	66	19	248%
2016-20	33	64	17	277%
2017-21	32	61	17	252%
2018-22	32	60	18	230%

Standardised Death Rate – Cancer U75

NI: Positive Change, Most Deprived: Positive Change, Least Deprived: Positive Change, Deprivation Gap: No Change

Years	Northern Ireland	Most Deprived	Least Deprived	Deprivation Gap
2014-18	146	202	120	69%
2015-19	142	196	115	71%
2016-20	140	193	113	70%
2017-21	136	188	109	74%
2018-22	133	183	105	74%

Standardised Death Rate – All Cause U75

NI: No Change, Most Deprived: No Change, Least Deprived: No Change, Deprivation Gap: No Change

Years	Northern Ireland	Most Deprived	Least Deprived	Deprivation Gap
2018	356	537	251	114%
2019	345	536	242	121%
2020	377	585	271	116%
2021	379	593	246	141%
2022	358	531	245	117%

Major Diseases

There were no changes in any of the most-least deprived gaps for indicators of major diseases. All admission indicators for major diseases saw decreases in rates within the most and least deprived areas, as well as regionally, while antihypertensive and statin prescription rates increased in all areas. The largest gaps observed remain for admissions due to respiratory diseases; particularly among those aged under 75 years, where the rate in the most deprived areas more than double the rate in the least deprived areas. The death rate due to COVID-19 in 2022 in the most deprived areas was one and a half times that in the least deprived areas.

Health Outcome Indicators

Standardised Prescription Rate – Antihypertensive

NI: Negative Change, Most Deprived: Negative Change, Least Deprived: Negative Change, Deprivation Gap: No Change

Years	Northern Ireland	Most Deprived	Least Deprived	Deprivation Gap
2018	216	246	199	24%
2019	214	239	197	21%
2020	223	253	204	24%
2021	225	255	206	24%
2022	228	258	210	23%

Standardised Prescription Rate – Statin

NI: Negative Change, Most Deprived: Negative Change, Least Deprived: Negative Change, Deprivation Gap: No Change

Years	Northern Ireland	Most Deprived	Least Deprived	Deprivation Gap
2018	166	194	146	33%
2019	167	196	147	34%
2020	164	194	144	35%
2021	167	195	146	34%
2022	171	199	150	32%

Standardised Incidence Rate - Cancer

NI: No Change, Most Deprived: No Change, Least Deprived: No Change, Deprivation Gap: No Change

Years	Northern Ireland	Most Deprived	Least Deprived	Deprivation Gap
2010-16	599	684	562	22%
2011-17	602	689	565	22%
2012-18	602	689	566	22%
2013-19	602	686	575	19%
2014-20	593	679	562	21%

Standardised Death Rate – COVID-19⁹

Years	Northern Ireland	Most Deprived	Least Deprived	Deprivation Gap
2020	101	124	98	27%
2021	111	145	76	90%
2022	39	49	33	50%

Service-based Indicators

Standardised Admission Rate – Circulatory^{10 11}

NI: Decrease, Most Deprived: Decrease, Least Deprived: Decrease, Deprivation Gap: No Change

Years	Northern Ireland	Most Deprived	Least Deprived	Deprivation Gap
2016/17-2018/19	1,986	2,164	1,742	24%
2017/18-2019/20	1,942	2,089	1,712	22%
2018/19-2020/21	1,721	1,839	1,528	20%
2019/20-2021/22	1,565	1,657	1,388	19%
2020/21-2022/23	1,441	1,535	1,284	20%

Standardised Admission Rate – Circulatory U75^{10 11}

NI: Decrease, Most Deprived: Decrease, Least Deprived: Decrease, Deprivation Gap: No Change

Years	Northern Ireland	Most Deprived	Least Deprived	Deprivation Gap
2016/17-2018/19	1,371	1,583	1,173	35%
2017/18-2019/20	1,341	1,529	1,144	34%
2018/19-2020/21	1,180	1,347	1,002	34%
2019/20-2021/22	1,061	1,205	892	35%
2020/21-2022/23	971	1,112	819	36%

Standardised Admission Rate – Respiratory^{10 12}

NI: Decrease, Most Deprived: Decrease, Least Deprived: Decrease, Deprivation Gap: No Change

Years	Northern Ireland	Most Deprived	Least Deprived	Deprivation Gap
2016/17-2018/19	2,017	2,819	1,452	94%
2017/18-2019/20	1,964	2,765	1,418	95%
2018/19-2020/21	1,553	2,234	1,108	102%
2019/20-2021/22	1,301	1,880	927	103%
2020/21-2022/23	1,162	1,671	850	97%

⁹ As COVID-19 is an emergent disease, data is only available for 2020 to 2022, therefore only a limited time series is available. The 'Standardised Death Rate – COVID-19' is based on deaths due to COVID-19 that have been registered with the General Register Office (GRO). It does not include deaths reported to the PHA where the deceased has had a positive test for COVID-19 and died within 28 days, where subsequently COVID-19 was not registered on the death certificate as the cause of death.

¹⁰ It should be noted that due to the impact of the COVID-19 pandemic on hospital services any changes from 2020/21 should be interpreted with caution.

¹¹ Circulatory admissions have been identified based on primary diagnosis field only.

¹² Respiratory admissions have been identified based on primary diagnosis field only.

Years	Northern Ireland	Most Deprived	Least Deprived	Deprivation Gap
2016/17-2018/19	1,462	2,142	1,006	113%
2017/18-2019/20	1,425	2,090	990	111%
2018/19-2020/21	1,112	1,666	753	121%
2019/20-2021/22	927	1,393	627	122%
2020/21-2022/23	826	1,247	570	119%

¹³ It should be noted that due to the impact of the COVID-19 pandemic on hospital services any changes from 2020/21 should be interpreted with caution.

¹⁴ Respiratory admissions have been identified based on primary diagnosis field only.

Hospital Activity

Across all hospital admission and attendance indicators, activity decreased over the analysed period in NI and the most and least deprived areas, and there were no notable changes observed in the inequality gaps, with the exception of the emergency admission rate where the most-least deprived gap narrowed slightly. The largest gaps occurred with emergency attendances and emergency care admissions.

Standardised Admission Rate – All Admissions^{15,16} NI: Decrease, Most Deprived: Decrease, Least Deprived: Decrease, Deprivation Gap: No Change

Years	Northern Ireland	Most Deprived	Least Deprived	Deprivation Gap
2018/19	23,395	27,762	19,863	40%
2019/20	22,434	26,238	19,345	36%
2020/21	14,864	17,971	12,908	39%
2021/22	18,224	21,852	15,738	39%
2022/23	20,042	23,441	17,328	35%

Standardised Admission Rate – Emergency Admissions^{15 16} NI: Decrease, Most Deprived: Decrease, Least Deprived: Decrease, Deprivation Gap: Narrowed

Years	Northern Ireland	Most Deprived	Least Deprived	Deprivation Gap
2018/19	8,885	11,216	6,832	64%
2019/20	8,749	11,001	6,847	61%
2020/21	6,892	8,799	5,422	62%
2021/22	7,715	9,755	6,025	62%
2022/23	7,651	9,455	6,214	52%

Standardised Attendance Rate – Emergency Care¹⁶ NI: Decrease, Most Deprived: Decrease, Least Deprived: Decrease, Deprivation Gap: No Change

Years	Northern Ireland	Most Deprived	Least Deprived	Deprivation Gap
2018/19	43,213	53,893	34,377	57%
2019/20	42,438	52,747	33,928	55%
2020/21	31,794	40,110	25,176	59%
2021/22	38,120	47,832	30,492	57%
2022/23	39,391	49,116	31,819	54%

¹⁵ Includes all admissions under the 'Acute Programme of Care' only.

¹⁶ It should be noted that due to the impact of the COVID-19 pandemic on hospital services any changes from 2020/21 should be interpreted with caution.

Standardised Admission Rate – Elective Inpatient Admissions^{17 18}

NI: Decrease, Most Deprived: Decrease, Least Deprived: Decrease, Deprivation Gap: No Change

Years	Northern Ireland	Most Deprived	Least Deprived	Deprivation Gap
2018/19	2,301	2,676	1,972	36%
2019/20	2,295	2,596	2,029	28%
2020/21	1,190	1,352	1,065	27%
2021/22	1,471	1,637	1,357	21%
2022/23	1,759	1,967	1,540	28%

Standardised Admission Rate – Day Case Admissions^{17 18}

NI: Decrease, Most Deprived: Decrease, Least Deprived: Decrease, Deprivation Gap: No Change

Years	Northern Ireland	Most Deprived	Least Deprived	Deprivation Gap
2018/19	11,049	12,608	10,149	24%
2019/20	10,321	11,409	9,608	19%
2020/21	5,714	6,663	5,475	22%
2021/22	7,983	9,357	7,444	26%
2022/23	9,590	10,885	8,696	25%

¹⁷ Includes all admissions under the 'Acute Programme of Care' only.

¹⁸ It should be noted that due to the impact of the COVID-19 pandemic on hospital services any changes from 2020/21 should be interpreted with caution.

Mental Health

Large inequality gaps continue to exist for mental health indicators however, there was a narrowing of the inequality gap in self-harm admissions, with a decrease in rates regionally and in the most and least deprived areas, but the rate in the most deprived areas remained around two and a half times the rate in the least deprived areas. It should be noted however that due to the impact of the COVID-19 pandemic on hospital services, changes should be interpreted with caution. Prescription rates for mood and anxiety disorders showed a negative change over the period with rates increasing regionally and in the most and least deprived areas. In 2020-22 the suicide mortality rate in the most deprived areas was more than two and a half times the rate observed in the least deprived areas.

Health Outcome Indicators

Crude Death Rate – Suicide¹⁹

NI: No Change, Most Deprived: No Change, Least Deprived: No Change, Deprivation Gap: No Change

Years	Northern Ireland	Most Deprived	Least Deprived	Deprivation Gap
2016-18	11.2	19.4	7.1	174%
2017-19	11.3	17.9	7.9	128%
2018-20	11.6	17.9	9.0	98%
2019-21	11.6	18.5	8.6	115%
2020-22	11.5	19.0	7.0	170%

Standardised Prescription Rate – Mood & Anxiety

NI: Negative Change, Most Deprived: Negative Change, Least Deprived: Negative Change, Deprivation Gap: No Change

Years	Northern Ireland	Most Deprived	Least Deprived	Deprivation Gap
2018	206	277	166	68%
2019	217	291	175	66%
2020	214	289	173	67%
2021	221	296	178	66%
2022	225	301	182	66%

¹⁹ It is important to note that registration-based figures build in delays arising from system wide processes which can drive annual fluctuations in the series. Events such as infant death, drug related deaths or suicide are usually referred to a coroner and this legal process can take some time.

Service-based Indicators

Standardised Admission Rate – Self-Harm^{20 21}

NI: Decrease, Most Deprived: Decrease, Least Deprived: Decrease, Deprivation Gap: Narrowed

Years	Northern Ireland	Most Deprived	Least Deprived	Deprivation Gap
2014/15-2018/19	136	250	74	240%
2015/16-2019/20	122	219	69	216%
2016/17-2020/21	111	190	66	189%
2017/18-2021/22	102	167	62	170%
2018/19-2022/23	91	147	58	154%

²⁰ It should be noted that due to the impact of the COVID-19 pandemic on hospital services any changes from 2020/21 should be interpreted with caution.

²¹ Self-harm admissions have been identified based on primary diagnosis field or any of the first 6 secondary diagnoses fields, and includes admissions under the 'Acute Programme of Care' only.

Alcohol, Smoking and Drugs

Alcohol, smoking and drugs related indicators continue to present some of the largest health inequalities in NI. The deprivation gap for drug misuse deaths widened over the analysed period and showed the largest inequality gap, where mortality in the most deprived areas was almost six times that of the least deprived. The gap for smoking attributable mortality also widened over the analysed period however rates improved in the most and least deprived areas. While inequality gaps in alcohol and drug related admissions narrowed over the period, the observed decreases in admission rates may be due to the impact of the COVID-19 pandemic on hospital services. Drug related, drug misuse and alcohol specific death rates all increased over the period both regionally and in the most and least deprived areas.²²

Health Outcome Indicators

Standardised Death Rate – Alcohol Specific

NI: Negative Change, Most Deprived: Negative Change, Least Deprived: Negative Change, Deprivation Gap: No Change

Years	Northern Ireland	Most Deprived	Least Deprived	Deprivation Gap
2014-18	15.8	31.5	7.6	315%
2015-19	17.0	34.7	8.3	318%
2016-20	17.6	36.6	9.0	306%
2017-21	18.1	38.0	8.6	342%
2018-22	18.5	39.3	9.1	334%

Standardised Death Rate – Smoking Attributable Causes

NI: Positive Change, Most Deprived: Positive Change, Least Deprived: Positive Change, Deprivation Gap: Widened

Years	Northern Ireland	Most Deprived	Least Deprived	Deprivation Gap
2014-18	249	363	186	95%
2015-19	242	354	180	97%
2016-20	234	342	173	97%
2017-21	224	328	165	99%
2018-22	212	315	157	101%

²² Please note that some observations may be due to changes in drug misuse behaviours among the population. For more information see [Appendix E: Technical Notes & Definitions](#).

Standardised Incidence Rate – Lung Cancer

NI: No Change, Most Deprived: No Change, Least Deprived: No Change, Deprivation Gap: No Change

Years	Northern Ireland	Most Deprived	Least Deprived	Deprivation Gap
2010-16	83	136	55	150%
2011-17	84	141	54	160%
2012-18	84	142	54	163%
2013-19	84	141	56	150%
2014-20	83	140	55	155%

Standardised Death Rate – Lung Cancer

NI: Positive Change, Most Deprived: Positive Change, Least Deprived: No Change, Deprivation Gap: Narrowed

Years	Northern Ireland	Most Deprived	Least Deprived	Deprivation Gap
2014-18	66	112	43	162%
2015-19	65	109	42	160%
2016-20	64	108	43	151%
2017-21	62	105	42	152%
2018-22	61	103	41	153%

Standardised Death Rate – Drug Related Causes²³

NI: Negative Change, Most Deprived: Negative Change, Least Deprived: Negative Change, Deprivation Gap: No Change

Years	Northern Ireland	Most Deprived	Least Deprived	Deprivation Gap
2014-18	7.7	17.9	3.7	386%
2015-19	8.5	20.3	4.1	399%
2016-20	9.3	21.4	4.6	363%
2017-21	10.2	22.6	5.6	307%
2018-22	10.4	23.4	4.7	392%

Standardised Death Rate – Drug Misuse²³

NI: Negative Change, Most Deprived: Negative Change, Least Deprived: No Change, Deprivation Gap: Widened

Years	Northern Ireland	Most Deprived	Least Deprived	Deprivation Gap
2014-18	6.3	14.7	2.8	425%
2015-19	7.1	17.0	3.0	461%
2016-20	7.8	18.1	3.5	412%
2017-21	8.5	19.8	3.7	441%
2018-22	8.7	20.0	3.4	494%

²³ It is important to note that registration-based figures build in delays arising from system wide processes which can drive annual fluctuations in the series. Events such as infant death, drug related deaths or suicide are usually referred to a coroner and this legal process can take some time.

Service-based Indicators

Standardised Admission Rate – Alcohol Related Causes^{24 25}

NI: Decrease, Most Deprived: Decrease, Least Deprived: Decrease, Deprivation Gap: Narrowed

Years	Northern Ireland	Most Deprived	Least Deprived	Deprivation Gap
2016/17-2018/19	669	1,400	350	300%
2017/18-2019/20	657	1,347	357	277%
2018/19-2020/21	609	1,243	331	276%
2019/20-2021/22	563	1,145	308	272%
2020/21-2022/23	517	1,026	294	250%

Standardised Admission Rate – Drug Related Causes^{24 25}

NI: Decrease, Most Deprived: Decrease, Least Deprived: Decrease, Deprivation Gap: Narrowed

Years	Northern Ireland	Most Deprived	Least Deprived	Deprivation Gap
2016/17-2018/19	190	372	103	261%
2017/18-2019/20	189	349	103	240%
2018/19-2020/21	178	322	98	227%
2019/20-2021/22	161	291	92	216%
2020/21-2022/23	136	252	80	213%

²⁴ It should be noted that due to the impact of the COVID-19 pandemic on hospital services any changes from 2020/21 should be interpreted with caution.

²⁵ Alcohol and drug related admissions have been identified based on primary diagnosis field or any of the first 6 secondary diagnoses fields and relates to admissions under the 'Acute Programme of Care' only.

Pregnancy & Early Years

Inequality gaps for smoking in pregnancy and teenage birth rate remained similar over the analysed period. While the gaps for the infant mortality rate and low birth weight widened slightly, the gap for breastfeeding on discharge narrowed due to an improvement in the rate in the most deprived areas while the rate in the least deprived areas saw no notable change. While improvements were observed regionally and in the most and least deprived areas with the under 20 teenage birth rate, in addition to regionally and in the most deprived areas for the proportion of mothers reporting smoking, the relative inequality gaps remain very large; with rates in the most deprived areas being over five times that in the least deprived.

Infant Mortality Rate²⁶

NI: No Change, Most Deprived: Negative Change, Least Deprived: No Change, Deprivation Gap: Widened

Years	Northern Ireland	Most Deprived	Least Deprived	Deprivation Gap
2014-18	4.5	4.8	4.3	12%
2015-19	4.6	5.3	3.7	45%
2016-20	4.4	5.2	3.7	40%
2017-21	4.3	5.1	4.0	28%
2018-22	4.4	5.3	4.2	26%

Smoking During Pregnancy

NI: Positive Change, Most Deprived: Positive Change, Least Deprived: No Change, Deprivation Gap: No Change

Years	Northern Ireland	Most Deprived	Least Deprived	Deprivation Gap
2018	13.2%	24.6%	4.4%	458%
2019	13.0%	23.4%	5.3%	340%
2020	13.0%	23.7%	5.0%	374%
2021	11.1%	21.3%	3.9%	440%
2022	10.8%	21.1%	4.1%	410%

Teenage Birth Rate U20

NI: Positive Change, Most Deprived: Positive Change, Least Deprived: Positive Change, Deprivation Gap: No Change

Years	Northern Ireland	Most Deprived	Least Deprived	Deprivation Gap
2018	8.4	17.1	3.4	402%
2019	8.1	17.3	2.8	514%
2020	6.2	13.4	1.9	603%
2021	5.9	11.0	2.4	360%
2022	5.3	10.2	2.0	421%

²⁶ As the underlying rates are relatively low (typically below 6 deaths per 1,000 live births), small annual changes can have a large impact on the resulting inequality gap. It is important to note that registration-based figures build in delays arising from system wide processes which can drive annual fluctuations in the series. Events such as infant death, drug related deaths or suicide are usually referred to a coroner and this legal process can take some time.

Breastfeeding on Discharge

NI: Positive Change, Most Deprived: Positive Change, Least Deprived: No Change, Deprivation Gap: Narrowed

Years	Northern Ireland	Most Deprived	Least Deprived	Deprivation Gap
2018	49%	34%	65%	48%
2019	50%	36%	66%	46%
2020	50%	35%	67%	48%
2021	51%	36%	68%	47%
2022	51%	38%	66%	42%

Low Birth Weight

NI: No Change, Most Deprived: No Change, Least Deprived: No Change, Deprivation Gap: Widened

Years	Northern Ireland	Most Deprived	Least Deprived	Deprivation Gap
2018	6.3%	7.4%	5.4%	38%
2019	6.1%	7.8%	5.4%	44%
2020	5.6%	7.5%	5.0%	50%
2021	5.9%	7.2%	4.6%	55%
2022	6.0%	8.0%	5.2%	54%

Healthy Birth Weight

NI: No Change, Most Deprived: No Change, Least Deprived: No Change, Deprivation Gap: No Change

Years	Northern Ireland	Most Deprived	Least Deprived	Deprivation Gap
2017	91%	91%	91%	0.6%
2018	91%	90%	92%	2.6%
2019	91%	89%	92%	2.8%
2020	91%	89%	91%	2.6%
2021	91%	90%	91%	1.1%

Small for Gestational Age²⁷

Years	Northern Ireland	Most Deprived	Least Deprived	Deprivation Gap
2019	10.3%	14.6%	8.2%	78%
2020	11.3%	14.4%	8.7%	65%
2021	10.7%	14.2%	8.6%	65%
2022	10.9%	13.7%	8.7%	56%

²⁷ No assessment of the inequality gap trend will be made until 5 years of data are available. Data is only presented for 2019 to 2022, due to insufficient recording levels prior to 2019. For further detail, please see [Appendix E: Technical Notes & Definitions](#).

Diet & Dental Health

In 2022/23 the percentage of primary 1 pupils in the most deprived areas affected by obesity was more than double the proportion in the least deprived areas. The inequality gap in year 8 pupils affected by obesity was slightly lower, with the proportion in the most deprived areas almost double that in the least deprived areas. The inequality gaps for most of the dental indicators showed no notable change over the analysed period and the dental registration rate showed no inequality gap between the most and least deprived areas for all ages.

Please note that BMI assessments are not available for 2019/20 and 2020/21 school years for Primary 1, and 2020/21 and 2021/22 school years for Year 8 BMI assessments, as the number of measurements recorded was negatively impacted by school closures due to the COVID-19 pandemic. An assessment of changes in the gap have therefore not been provided and caution should be taken with the assessments of change in the rates provided below.

In addition, Year 8 BMI data for 2018/19 and 2019/20 does not include measurements across all HSC Trusts therefore figures for these years should be treated with caution.

Health Outcome Indicators

Primary 1 BMI: Obese²⁸

NI: No Change, Most Deprived: No Change, Least Deprived: Positive Change

Years	Northern Ireland	Most Deprived	Least Deprived	Deprivation Gap
2018/19	5.5%	6.6%	4.2%	55%
2019/20				
2020/21				
2021/22	6.1%	8.7%	3.8%	129%
2022/23	5.1%	7.0%	3.0%	130%

Primary 1 BMI: Overweight or Obese²⁸

NI: No Change, Most Deprived: No Change, Least Deprived: No Change

Years	Northern Ireland	Most Deprived	Least Deprived	Deprivation Gap
2018/19	22%	25%	18%	36%
2019/20				
2020/21				
2021/22	22%	26%	18%	48%
2022/23	20%	24%	17%	40%

²⁸ Assessment of change in the inequality gap has not been provided due to limited data series as a result of the pandemic. See page 94 for further information.

Year 8 BMI: Obese²⁹

NI: No Change, Most Deprived: Positive Change, Least Deprived: No Change

Years	Northern Ireland	Most Deprived	Least Deprived	Deprivation Gap
2018/19	5.8%	8.7%	3.8%	128%
2019/20	6.0%	8.7%	3.8%	127%
2020/21				
2021/22				
2022/23	5.6%	7.5%	3.9%	94%

Year 8 BMI: Overweight or Obese²⁹

NI: No Change, Most Deprived: No Change, Least Deprived: No Change

Years	Northern Ireland	Most Deprived	Least Deprived	Deprivation Gap
2018/19	27%	34%	22%	54%
2019/20	26%	33%	22%	52%
2020/21				
2021/22				
2022/23	26%	32%	22%	47%

Service-based Indicators**Standardised Dental Filling Rate^{30 31}**

NI: Decrease, Most Deprived: Decrease, Least Deprived: Decrease, Deprivation Gap: No Change

Years	Northern Ireland	Most Deprived	Least Deprived	Deprivation Gap
2018/19	50,305	60,924	41,162	48%
2019/20	47,365	57,619	38,391	50%
2020/21	13,133	13,939	11,425	22%
2021/22	24,797	27,300	20,957	30%
2022/23	35,777	42,035	29,142	44%

Standardised Dental Filling Rate – U18^{30 31}

NI: Decrease, Most Deprived: Decrease, Least Deprived: Decrease, Deprivation Gap: No Change

Years	Northern Ireland	Most Deprived	Least Deprived	Deprivation Gap
2018/19	45,438	55,017	33,535	64%
2019/20	42,475	50,552	32,495	56%
2020/21	10,520	11,057	8,721	27%
2021/22	20,915	22,708	16,377	39%
2022/23	33,276	39,406	25,170	57%

²⁹ Assessment of change in the inequality gap has not been provided due to limited data series as a result of the pandemic. See page 94 for further information.

³⁰ Due to the major impact of the COVID-19 pandemic on dental services any changes in 2020/21 and 2021/22 should be interpreted with caution. It is also important to note that data does not include those who have paid privately for treatment.

³¹ Figures relate to the total procedures' individuals have had carried out within the given period (i.e., includes multiple procedures for an individual where applicable). For 'number of individuals that have undergone at least one procedure' see 'additional indicators' section.

Standardised Dental Extraction Rate^{32 33} NI: Decrease, Most Deprived: Decrease, Least Deprived: Decrease, Deprivation Gap: No Change

Years	Northern Ireland	Most Deprived	Least Deprived	Deprivation Gap
2018/19	8,969	12,950	6,366	103%
2019/20	8,563	12,430	6,077	105%
2020/21	4,688	6,534	3,331	96%
2021/22	6,711	9,766	4,665	109%
2022/23	7,647	11,056	5,309	108%

Standardised Dental Extraction Rate - U18^{32 33} NI: Decrease, Most Deprived: Decrease, Least Deprived: Decrease, Deprivation Gap: Narrowed

Years	Northern Ireland	Most Deprived	Least Deprived	Deprivation Gap
2018/19	7,278	7,948	6,844	16%
2019/20	7,164	7,637	6,864	11%
2020/21	3,369	3,285	3,267	1%
2021/22	4,991	5,223	4,706	11%
2022/23	6,087	6,267	5,852	7%

Standardised Dental Crowning Rate^{32 33} NI: Decrease, Most Deprived: Decrease, Least Deprived: Decrease, Deprivation Gap: No Change

Years	Northern Ireland	Most Deprived	Least Deprived	Deprivation Gap
2018/19	2,081	2,642	1,719	54%
2019/20	1,873	2,446	1,585	54%
2020/21	411	443	412	8%
2021/22	702	806	679	19%
2022/23	995	1,228	908	35%

Standardised Dental Registration Rate NI: Increase, Most Deprived: Increase, Least Deprived: Increase, Deprivation Gap: Narrowed

Years	Northern Ireland	Most Deprived	Least Deprived	Deprivation Gap
2018/19	63,626	62,735	64,251	2.4%
2019/20	63,560	62,450	64,240	2.8%
2020/21	65,075	64,070	65,345	2.0%
2021/22	67,252	66,399	67,134	1.1%
2022/23	69,829	69,380	69,259	-0.2%

³² Due to the major impact of the COVID-19 pandemic on dental services any changes in 2020/21 and 2021/22 should be interpreted with caution. It is also important to note that data does not include those who have paid privately for treatment.

³³ Figures relate to the total procedures' individuals have had carried out within the given period (i.e., includes multiple procedures for an individual where applicable). For 'number of individuals that have undergone at least one procedure' see 'additional indicators' section.

Standardised Dental Registration Rate (U18)NI: No Change, Most Deprived: No Change, Least Deprived:
No Change, Deprivation Gap: No Change

Years	Northern Ireland	Most Deprived	Least Deprived	Deprivation Gap
2018/19	76,227	71,252	79,898	10.8%
2019/20	76,286	71,066	80,264	11.5%
2020/21	73,865	68,886	77,494	11.1%
2021/22	73,561	68,559	76,965	10.9%
2022/23	75,761	71,448	79,360	10.0%

Sub-regional Health Inequalities

There are two aspects to the analysis of sub-regional health inequalities in this section. The analysis for each sub-regional area includes both the difference in health outcomes between the **Trust or LGD and the regional (NI) average** or to the **20% most deprived areas within an area and the area's average**.

Each chapter is a summary of findings only. For a full assessment of the HSC Trust and LGD figures, including a range of indicators that are also available at District Electoral Area (DEA), see downloadable tables at:

<https://www.health-ni.gov.uk/articles/health-inequalities-statistics>

Each chapter contains a comparison of the area's health outcomes against the regional average. This includes:

- Detail of life expectancy at birth in the Trust/LGD and how it compares with the NI average.
- A table which presents how each health outcome indicator for the Trust/LGD compares with the NI average. Gaps can be significantly better or worse than the NI average, or, where there is no statistically significant difference, they are categorised as 'similar to'. Male and female life expectancy at birth and at age 65 gaps are not included in this chart as they are represented in years.
- Outcomes relating to **service-based indicators**, including hospital admissions, are **not presented** here as it is not possible to ascertain whether a Trust-NI gap in utilisation of services fully constitutes a difference in health outcomes due to potentially different service provision profiles across different geographic areas. For further information see 'Notes for users' Page 9.

There is also a comparison of the Inequality Gap between the area average and its most deprived areas in respect of:

- Life Expectancy: Analysis of the male and female life expectancy gap within the area.
- Largest Inequality Gaps: The five largest inequality gaps in the area. Gaps that are not statistically significant are not included in this ranking.
- Outcomes relating to **service-based indicators**, including hospital admissions, **are included** here. For further information see 'Notes for users' Page 9.

COMPARISON OF TRUST HEALTH OUTCOMES AGAINST THE REGIONAL AVERAGE

The following Trusts had a majority of health outcomes that were better than the NI average:

- South Eastern HSC Trust

The following Trusts had a majority of health outcomes that were worse than the NI average:

- Belfast HSC Trust

Note: Outcomes relating to service-based indicators, including hospital admissions, have not been presented here due to difficulties in ascertaining whether any differences in rates are due to differences in demand (i.e., health of the population), or, because of access difficulties (For further information see ‘Notes for users’ Page 9).

Trust	Better Than	Similar To	Worse Than
Belfast	0	8	26
Northern	14	20	0
South Eastern	18	16	0
Southern	16	17	1
Western	1	27	6

Largest Deprivation Inequality Gaps in Each HSC Trust Area

The table below indicates the five largest deprivation inequality gaps in each Health & Social Care Trust (HSCT).^{34 35}

Belfast HSCT	Drug Misuse Death Rate (128%)	Drug Related Death Rate (117%)	Alcohol Related Admission Rate (100%)	Under 20 Teenage Birth Rate (99%)	Under 75 Respiratory Death Rate (99%)
Northern HSCT	Drug Related Death Rate (133%)	Drug Misuse Death Rate (129%)	Drug Related Admissions (108%)	Smoking during Pregnancy (101%)	Self-Harm Admission Rate (96%)
South Eastern HSCT	Drug Misuse Death Rate (112%)	Drug Related Death Rate (110%)	Under 20 Teenage Birth Rate (102%)	Alcohol Specific Death Rate (99%)	Drug Related Admission Rate (99%)
Southern HSCT	Alcohol Specific Death Rate (101%)	Alcohol Related Admission Rate (96%)	Under 20 Teenage Birth Rate (89%)	Drug Related Admission Rate (88%)	Smoking During Pregnancy (82%)
Western HSCT	Drug Misuse Death Rate (145%)	Drug Related Death Rate (138%)	Alcohol Related Admission Rate (115%)	Alcohol Specific Death Rate (107%)	Under 20 Teenage Birth Rate (100%)

³⁴ Includes all indicators including outcomes relating to service-based indicators. For further information see ‘Notes for users’, Page 9

³⁵ Excludes gaps that were not statistically significant.

Belfast

Health & Social Care Trust (HSCT)

Comparison of Health Outcomes with NI³⁶

26 Health Outcomes were worse than the NI Average

8 Health Outcomes were similar to the NI Average

Of the 34 health outcome indicators analysed for Belfast Trust residents, 26 were worse than the NI average, eight were similar to the NI average, and none of the health outcomes analysed were better than the NI average.

Life Expectancy:

In the Belfast Trust, male life expectancy (76.3 years) was 2.1 years less than the NI average (78.4 years) and female life expectancy (80.8 years) was 1.4 years less than the average (82.3 years).

Gaps in Health Outcomes between Belfast Trust and NI³⁷

Life expectancy at birth and at age 65 gaps included in the figures above have not been displayed in the table below as they are represented in years.

Indicator	Gap	Comparison to NI
Standardised Death Rate - Drug Misuse	80%	Worse Than
Standardised Death Rate - Drug Related Causes	75%	Worse Than
Teenage Birth Rate U20	56%	Worse Than
Standardised Death Rate - Respiratory U75	45%	Worse Than
Standardised Death Rate - Alcohol Specific	42%	Worse Than
Smoking During Pregnancy	41%	Worse Than
Crude Suicide Rate	39%	Worse Than
Standardised Death Rate - Lung Cancer	39%	Worse Than
Standardised Death Rate - Preventable	35%	Worse Than
Standardised Incidence Rate - Lung Cancer	32%	Worse Than
Standardised Death Rate - Avoidable	31%	Worse Than
Standardised Death Rate - All Cause U75	26%	Worse Than
Potential Years of Life Lost	25%	Worse Than
Standardised Death Rate - Smoking Attributable Causes	23%	Worse Than
Standardised Death Rate - Cancer U75	22%	Worse Than
Standardised Death Rate - Treatable	22%	Worse Than
Standardised Death Rate - Circulatory U75	17%	Worse Than
Low Birth Weight	15%	Worse Than
Small for Gestational Age	13%	Similar To
Standardised Death Rate - All Deaths	12%	Worse Than
Standardised Prescription Rate - Mood & Anxiety	10%	Worse Than

³⁶ Outcomes relating to service-based indicators, including hospital admissions, have not been presented above as it is not possible to ascertain whether a Trust-NI gap in utilisation of services fully constitutes a difference in health outcomes due to potentially different service provision profiles across different geographic areas. For further information see 'Notes for users', Page 9.

³⁷ As confidence intervals vary across indicators, some gaps between the Trust and NI are not significant and have therefore been categorised as 'similar to', despite being larger than gaps for other indicators that have been identified as significant (i.e. worse than or better than).

Standardised Incidence Rate - Cancer	9%	Worse Than
Standardised Death Rate - COVID-19	5%	Similar To
Standardised Prescription Rate - Antihypertensive	4%	Worse Than
Standardised Prescription Rate - Statin	0%	Similar To
Breastfeeding on Discharge	-3%	Similar To
Year 8 BMI: Overweight or Obese	-3%	Similar To
Primary 1 BMI: Overweight or Obese	-4%	Similar To
Year 8 BMI: Obese	-4%	Similar To
Primary 1 BMI: Obese	-5%	Similar To

Comparison with Most Deprived Areas

Inequality gaps between the 20% most deprived areas within the Belfast Trust and the Trust average:

Life Expectancy

- Male life expectancy in the Trust's most deprived areas was 70.8 years, **5.5 years** less than the HSCT average (76.3 years).
- Female life expectancy in the Trust's most deprived areas was 77.0 years, **3.9 years** less than the HSCT average (80.8 years).

Largest Inequality Gaps³⁸

Across the 53 indicators analysed, the majority of outcomes were significantly worse for those residing in the 20% most deprived areas of Belfast Trust when compared with the Belfast Trust average. Exceptions include deaths due to covid, babies born small for gestational age, and primary 1 & year 8 overweight and/or affected by obesity, where differences were not statistically significant. The largest inequality gaps observed in the latest year were³⁹:

Drug Misuse Death Rate (128%)	Drug Related Death Rate (117%)	Alcohol Related Admission Rate (100%)	Under 20 Teenage Birth Rate (99%)	Under 75 Respiratory Death Rate (99%)
-------------------------------	--------------------------------	---------------------------------------	-----------------------------------	---------------------------------------

This is a summary of findings only. For a full assessment and all figures see downloadable tables at:

<https://www.health-ni.gov.uk/articles/health-inequalities-statistics>

³⁸ Includes all indicators including outcomes relating to service-based indicators. For further information see 'Notes for users', Page 9.

³⁹ Excludes gaps that were not statistically significant.

Northern

Health & Social Care Trust (HSCT)

Comparison with NI⁴⁰

20 Health Outcomes were similar to the NI Average

14 Health Outcomes were better than the NI Average

Of the 34 health outcome indicators analysed for Northern Trust residents, none were worse than the NI average, 20 were similar to the NI average, and 14 of the health outcomes analysed were better than the NI average.

Life Expectancy:

In the Northern Trust, male life expectancy (78.9 years) was 0.4 years higher than the NI average (78.4 years) and female life expectancy (82.8 years) was 0.5 years higher than the average (82.3 years).

Gaps in Health Outcomes between Northern Trust and NI⁴¹

Life expectancy gaps at birth and at age 65 included in the figures above have not been displayed in the table below as they are represented in years.

Indicator	Gap	Comparison to NI
Primary 1 BMI: Obese	15%	Similar To
Primary 1 BMI: Overweight or Obese	6%	Similar To
Smoking During Pregnancy	2%	Similar To
Year 8 BMI: Obese	1%	Similar To
Breastfeeding on Discharge	0%	Similar To
Standardised Prescription Rate - Antihypertensive	-1%	Better Than
Standardised Death Rate - Circulatory U75	-1%	Similar To
Standardised Prescription Rate - Mood & Anxiety	-2%	Better Than
Standardised Incidence Rate - Cancer	-2%	Better Than
Standardised Death Rate - All Deaths	-2%	Similar To
Standardised Prescription Rate - Statin	-4%	Better Than
Standardised Death Rate - Treatable	-4%	Similar To
Small for Gestational Age	-4%	Similar To
Standardised Death Rate - Smoking Attributable Causes	-4%	Similar To
Standardised Death Rate - Respiratory U75	-5%	Similar To
Standardised Death Rate - All Cause U75	-5%	Similar To
Standardised Death Rate - Cancer U75	-6%	Similar To
Year 8 BMI: Overweight or Obese	-6%	Similar To
Standardised Death Rate - Avoidable	-7%	Better Than
Teenage Birth Rate U20	-7%	Similar To
Low Birth Weight	-7%	Better Than
Potential Years of Life Lost	-8%	Better Than

⁴⁰ Outcomes relating to service-based indicators, including hospital admissions, have not been presented above as it is not possible to ascertain whether a Trust-NI gap in utilisation of services fully constitutes a difference in health outcomes due to potentially different service provision profiles across different geographic areas. For further information see 'Notes for users', Page 9.

⁴¹ As confidence intervals vary across indicators, some gaps between the Trust and NI are not significant and have therefore been categorised as 'similar to', despite being larger than gaps for other indicators that have been identified as significant (i.e. worse than or better than).

Standardised Incidence Rate - Lung Cancer	-8%	Better Than
Standardised Death Rate - Preventable	-8%	Better Than
Standardised Death Rate - Lung Cancer	-10%	Better Than
Standardised Death Rate - Drug Misuse	-15%	Similar To
Standardised Death Rate - COVID-19	-16%	Similar To
Standardised Death Rate - Drug Related Causes	-17%	Similar To
Standardised Death Rate - Alcohol Specific	-18%	Better Than
Crude Suicide Rate	-21%	Better Than

Comparison with Most Deprived Areas

Inequality gaps between the 20% most deprived areas within the Northern Trust and the Trust average:

Life Expectancy

- Male life expectancy in the Trust's most deprived areas was 75.0 years, **3.9 years** less than the HSCT average (78.9 years).
- Female life expectancy in the Trust's most deprived areas was 80.6 years, **2.2 years** less than the HSCT average (82.8 years).

Largest Inequality Gaps within Northern Trust⁴²

Across the 53 indicators analysed, the majority of outcomes were significantly worse for those residing in the 20% most deprived areas of Northern Trust when compared with the Northern Trust average. Exceptions include under 75 cancer mortality, babies born small for gestational age, and primary 1 & year 8 overweight and/or affected by obesity, where differences were not statistically significant. The largest inequality gaps observed in the latest year were⁴³:

Drug Related Death Rate (133%)	Drug Misuse Death Rate (129%)	Drug Related Admissions (108%)	Smoking during Pregnancy (101%)	Self-Harm Admission Rate (96%)
--------------------------------	-------------------------------	--------------------------------	---------------------------------	--------------------------------

This is a summary of findings only. For a full assessment and all figures see downloadable tables at:

<https://www.health-ni.gov.uk/articles/health-inequalities-statistic>

⁴² Includes all indicators including outcomes relating to service-based indicators. For further information see 'Notes for users', Page 9.

⁴³ Excludes gaps that were not statistically significant.

South Eastern

Health & Social Care Trust (HSCT)

Comparison with NI⁴⁴

16 Health Outcomes were similar to the NI Average

18 Health Outcomes were better than the NI Average

Of the 34 health outcome indicators analysed for South Eastern Trust residents, none were worse than the NI average, 16 were similar to the NI average, and 18 of the health outcomes analysed were better than the NI average.

Life Expectancy:

In the South Eastern Trust, male life expectancy (79.7 years) was 1.3 years higher than the NI average (78.4 years) and female life expectancy (82.7 years) was similar to the average (82.3 years).

Gaps in Health Outcomes between South Eastern Trust and NI⁴⁵

Life expectancy gaps at birth and at age 65 included in the figures above have not been displayed in the table below as they are represented in years.

Indicator	Gap	Comparison to NI
Teenage Birth Rate U20	7%	Similar To
Standardised Death Rate - COVID-19	3%	Similar To
Low Birth Weight	3%	Similar To
Standardised Prescription Rate - Antihypertensive	-1%	Better Than
Small for Gestational Age	-1%	Similar To
Primary 1 BMI: Overweight or Obese	-2%	Similar To
Year 8 BMI: Overweight or Obese	-2%	Similar To
Standardised Prescription Rate - Mood & Anxiety	-3%	Better Than
Standardised Incidence Rate - Cancer	-4%	Better Than
Standardised Death Rate - Cancer U75	-5%	Similar To
Crude Suicide Rate	-5%	Similar To
Breastfeeding on Discharge	-5%	Similar To
Standardised Death Rate - All Deaths	-6%	Better Than
Standardised Prescription Rate - Statin	-8%	Better Than
Standardised Death Rate - Treatable	-8%	Better Than
Potential Years of Life Lost	-10%	Better Than
Standardised Death Rate - All Cause U75	-10%	Better Than
Standardised Death Rate - Smoking Attributable Causes	-10%	Better Than
Primary 1 BMI: Obese	-11%	Similar To
Standardised Death Rate - Drug Related Causes	-11%	Similar To
Standardised Death Rate - Circulatory U75	-11%	Similar To

⁴⁴ Outcomes relating to service-based indicators, including hospital admissions, have not been presented above as it is not possible to ascertain whether a Trust-NI gap in utilisation of services fully constitutes a difference in health outcomes due to potentially different service provision profiles across different geographic areas. For further information see 'Notes for users', Page 9.

⁴⁵ As confidence intervals vary across indicators, some gaps between the Trust and NI are not significant and have therefore been categorised as 'similar to', despite being larger than gaps for other indicators that have been identified as significant (i.e. worse than or better than).

Standardised Death Rate - Drug Misuse	-12%	Similar To
Standardised Death Rate - Avoidable	-13%	Better Than
Standardised Death Rate - Lung Cancer	-13%	Better Than
Smoking During Pregnancy	-14%	Better Than
Standardised Death Rate - Preventable	-15%	Better Than
Standardised Incidence Rate - Lung Cancer	-18%	Better Than
Standardised Death Rate - Respiratory U75	-18%	Similar To
Year 8 BMI: Obese	-20%	Similar To
Standardised Death Rate - Alcohol Specific	-25%	Better Than

Comparison with Most Deprived Areas

Inequality gaps between the 20% most deprived areas within the South Eastern Trust and the Trust average:

Life Expectancy

- Male life expectancy in the Trust's most deprived areas was 75.7 years, **4.0 years** less than the HSCT average (79.7 years).
- Female life expectancy in the Trust's most deprived areas was 80.2 years, **2.5 years** less than the HSCT average (82.7 years).

Largest Inequality Gaps within South Eastern Trust⁴⁶

Across the 53 indicators analysed, the majority of outcomes were significantly worse for those residing in the 20% most deprived areas of South Eastern Trust when compared with the South Eastern Trust average. Exceptions include low birth weight, babies born small for gestational age, and primary 1 & year 8 overweight and/or affected by obesity, where differences were not statistically significant. The largest inequality gaps observed in the latest year were⁴⁷:

Drug Misuse Death Rate (112%)	Drug Related Death Rate (110%)	Under 20 Teenage Birth Rate (102%)	Alcohol Specific Death Rate (99%)	Drug Related Admission Rate (99%)
-------------------------------	--------------------------------	------------------------------------	-----------------------------------	-----------------------------------

This is a summary of findings only. For a full assessment and all figures see downloadable tables at:
<https://www.health-ni.gov.uk/articles/health-inequalities-statistics>

⁴⁶ Includes all indicators including outcomes relating to service-based indicators. For further information see 'Notes for users', Page 9.

⁴⁷ Excludes gaps that were not statistically significant.

Southern

Health & Social Care Trust (HSCT)

Comparison with NI⁴⁸

1 Health Outcome was worse than the NI Average

17 Health Outcomes were similar to the NI Average

16 Health Outcomes were better than the NI Average

Of the 34 health outcome indicators analysed for Southern Trust residents, one was worse than the NI average, 17 were similar to the NI average, and 16 of the health outcomes analysed were better than the NI average.

Life Expectancy:

In the Southern Trust, male life expectancy (78.7 years) was similar to the NI average (78.4 years) and female life expectancy (82.7 years) was 0.5 years higher than the average (82.3 years).

Gaps in Health Outcomes between Southern Trust and NI⁴⁹

Life expectancy gaps at birth and at age 65 included in the figures above have not been displayed in the table below as they are represented in years.

Indicator	Gap	Comparison to NI
Year 8 BMI: Obese	8%	Similar To
Standardised Death Rate - COVID-19	4%	Similar To
Standardised Prescription Rate - Statin	4%	Worse Than
Year 8 BMI: Overweight or Obese	2%	Similar To
Standardised Prescription Rate - Antihypertensive	-1%	Similar To
Breastfeeding on Discharge	-2%	Similar To
Primary 1 BMI: Overweight or Obese	-2%	Similar To
Standardised Incidence Rate - Cancer	-2%	Better Than
Standardised Death Rate - All Deaths	-3%	Better Than
Potential Years of Life Lost	-4%	Similar To
Standardised Death Rate - Circulatory U75	-5%	Similar To
Standardised Death Rate - Cancer U75	-5%	Similar To
Standardised Incidence Rate - Lung Cancer	-5%	Better Than
Low Birth Weight	-6%	Similar To
Standardised Death Rate - Treatable	-6%	Similar To
Standardised Death Rate - Alcohol Specific	-6%	Similar To
Primary 1 BMI: Obese	-6%	Similar To
Standardised Prescription Rate - Mood & Anxiety	-6%	Better Than
Standardised Death Rate - All Cause U75	-7%	Better Than
Standardised Death Rate - Avoidable	-8%	Better Than
Standardised Death Rate - Smoking Attributable Causes	-8%	Better Than

⁴⁸ Outcomes relating to service-based indicators, including hospital admissions, have not been presented above as it is not possible to ascertain whether a Trust-NI gap in utilisation of services fully constitutes a difference in health outcomes due to potentially different service provision profiles across different geographic areas. For further information see 'Notes for users', Page 9.

⁴⁹ As confidence intervals vary across indicators, some gaps between the Trust and NI are not significant and have therefore been categorised as 'similar to', despite being larger than gaps for other indicators that have been identified as significant (i.e. worse than or better than).

Standardised Death Rate - Preventable	-8%	Better Than
Standardised Death Rate - Lung Cancer	-10%	Better Than
Crude Suicide Rate	-10%	Similar To
Small for Gestational Age	-13%	Better Than
Standardised Death Rate - Respiratory U75	-19%	Better Than
Smoking During Pregnancy	-21%	Better Than
Teenage Birth Rate U20	-23%	Better Than
Standardised Death Rate - Drug Related Causes	-38%	Better Than
Standardised Death Rate - Drug Misuse	-41%	Better Than

Comparison with Most Deprived Areas

Inequality gaps between the 20% most deprived areas within the Southern Trust and the Trust average:

Life Expectancy

- Male life expectancy in the Trust's most deprived areas was 75.8 years, **2.9 years** less than the HSC average (78.7 years).
- Female life expectancy in the Trust's most deprived areas was 81.7 years, **1.0 years** less than the HSC average (82.7 years).

Largest Inequality Gaps within Southern Trust⁵⁰

Across the 53 indicators analysed, the majority of outcomes were significantly worse for those residing in the 20% most deprived areas of Southern Trust when compared with the Southern Trust average. Exceptions include under 75 cancer death rate, deaths due to covid, suicide rate, drug related and drugs misuse death rates, low birth weight, babies born small for gestational age, and primary 1 & year 8 overweight and/or affected by obesity, where differences were not statistically significant. The largest inequality gaps observed in the latest year were⁵¹:

Alcohol Specific Death Rate (101%)	Alcohol Related Admission Rate (96%)	Under 20 Teenage Birth Rate (89%)	Drug Related Admission Rate (88%)	Smoking During Pregnancy (82%)
------------------------------------	--------------------------------------	-----------------------------------	-----------------------------------	--------------------------------

This is a summary of findings only. For a full assessment and all figures see downloadable tables at:
<https://www.health-ni.gov.uk/articles/health-inequalities-statistics>

⁵⁰ Includes all indicators including outcomes relating to service-based indicators. For further information see 'Notes for users', Page 9.

⁵¹ Excludes gaps that were not statistically significant.

Western

Health & Social Care Trust (HSCT)

Comparison with NI⁵²

6 Health Outcome were worse than the NI Average

27 Health Outcomes were similar to the NI Average

1 Health Outcomes was better than the NI Average

Of the 34 health outcome indicators analysed for Western Trust residents, six were worse than the NI average, 27 were similar to the NI average, and one of the health outcomes analysed was better than the NI average.

Life Expectancy:

In the Western Trust, male life expectancy (78.2 years) was similar to the NI average (78.4 years) and female life expectancy (82.0 years) was also similar to the average (82.3 years).

Gaps in Health Outcomes between Western Trust and NI⁵³

Life expectancy gaps at birth and at age 65 included in the figures above have not been displayed in the table below as they are represented in years.

Indicator	Gap	Comparison to NI
Standardised Death Rate - Alcohol Specific	22%	Worse Than
Standardised Death Rate - COVID-19	14%	Similar To
Standardised Prescription Rate - Statin	12%	Worse Than
Breastfeeding on Discharge	12%	Worse Than
Small for Gestational Age	10%	Similar To
Year 8 BMI: Overweight or Obese	10%	Similar To
Standardised Incidence Rate - Lung Cancer	7%	Worse Than
Standardised Prescription Rate - Mood & Anxiety	7%	Worse Than
Standardised Death Rate - Smoking Attributable Causes	6%	Worse Than
Standardised Death Rate - Respiratory U75	6%	Similar To
Crude Suicide Rate	6%	Similar To
Year 8 BMI: Obese	6%	Similar To
Standardised Death Rate - Circulatory U75	4%	Similar To
Standardised Death Rate - Lung Cancer	4%	Similar To
Standardised Death Rate - Preventable	3%	Similar To
Primary 1 BMI: Obese	3%	Similar To
Standardised Death Rate - All Deaths	3%	Worse Than
Standardised Death Rate - Avoidable	3%	Similar To
Potential Years of Life Lost	2%	Similar To
Standardised Incidence Rate - Cancer	1%	Similar To
Standardised Death Rate - Treatable	1%	Similar To

⁵² Outcomes relating to service-based indicators, including hospital admissions, have not been presented above as it is not possible to ascertain whether a Trust-NI gap in utilisation of services fully constitutes a difference in health outcomes due to potentially different service provision profiles across different geographic areas. For further information see 'Notes for users', Page 9.

⁵³ As confidence intervals vary across indicators, some gaps between the Trust and NI are not significant and have therefore been categorised as 'similar to', despite being larger than gaps for other indicators that have been identified as significant (i.e. worse than or better than).

Standardised Death Rate - All Cause U75	1%	Similar To
Primary 1 BMI: Overweight or Obese	0%	Similar To
Standardised Prescription Rate - Antihypertensive	-1%	Similar To
Low Birth Weight	-2%	Similar To
Standardised Death Rate - Cancer U75	-2%	Similar To
Smoking During Pregnancy	-4%	Similar To
Standardised Death Rate - Drug Related Causes	-8%	Similar To
Standardised Death Rate - Drug Misuse	-15%	Similar To
Teenage Birth Rate U20	-30%	Better Than

Comparison with Most Deprived Areas

Inequality gaps between the 20% most deprived areas within the Western Trust and the Trust average:

Life Expectancy

- Male life expectancy in the Trust's most deprived areas was 73.2 years, **5.0 years** less than the HSC average (78.2 years).
- Female life expectancy in the Trust's most deprived areas was 79.3 years, **2.7 years** less than the HSC average (82.0 years).

Largest Inequality Gaps within Western Trust⁵⁴

Across the 53 indicators analysed, the majority of outcomes were significantly worse for those residing in the 20% most deprived areas of Western Trust when compared with the Western Trust average. Exceptions include deaths due to covid (where the rate was lower in the most deprived areas), and low birth weight, babies born small for gestational age, and primary 1 & year 8 overweight and/or affected by obesity, where differences were not statistically significant. The largest inequality gaps observed in the latest year were⁵⁵:

Drug Misuse Death Rate (145%)	Drug Related Death Rate (138%)	Alcohol Related Admission Rate (115%)	Alcohol Specific Death Rate (107%)	Under 20 Teenage Birth Rate (100%)
-------------------------------	--------------------------------	---------------------------------------	------------------------------------	------------------------------------

This is a summary of findings only. For a full assessment and all figures see downloadable tables at:
<https://www.health-ni.gov.uk/articles/health-inequalities-statistics>

⁵⁴ Includes all indicators including outcomes relating to service-based indicators. For further information see 'Notes for users', Page 9.

⁵⁵ Excludes gaps that were not statistically significant.

COMPARISON OF LGD HEALTH OUTCOMES AGAINST THE REGIONAL AVERAGE

The following LGDs had a majority of health outcomes that were better than the NI average:

- Ards & North Down LGD
- Lisburn & Castlereagh LGD
- Mid Ulster LGD

The following LGDs had a majority of health outcomes that were worse than the NI average:

- Belfast LGD

Note: Outcomes relating to service-based indicators, including hospital admissions, have not been presented here due to difficulties in ascertaining whether any differences in rates are due to differences in demand (i.e., health of the population), or, because of access difficulties (For further information see 'Notes for users' Page 9).

Local Government District	Better Than	Similar To	Worse Than
Antrim & Newtownabbey	2	30	0
Ards & North Down	21	11	0
Armagh City, Banbridge & Craigavon	12	20	0
Belfast	0	8	24
Causeway Coast & Glens	10	21	1
Derry City & Strabane	0	17	15
Fermanagh & Omagh	8	23	1
Lisburn & Castlereagh	22	10	0
Mid & East Antrim	3	27	2
Mid Ulster	17	14	1
Newry, Mourne & Down	7	24	1

Largest Deprivation Inequality Gaps in Each LGD

The table below indicates the five largest deprivation inequality gaps in each Local Government District (LGD).^{56 57}

Antrim & Newtownabbey LGD	Drug Related Death Rate (114%)	Alcohol Specific Death Rate (111%)	Self-Harm Admission Rate (93%)	Smoking during Pregnancy (85%)	Drug Related Admission Rate (76%)
Ards & North Down LGD	Drug Related Admission Rate (97%)	Smoking during Pregnancy (81%)	Alcohol Related Admission Rate (75%)	Preventable Death Rate (67%)	Self-Harm Admission Rate (64%)
Armagh City, Banbridge & Craigavon LGD	Under 20 Teenage Birth Rate (134%)	Smoking During Pregnancy (111%)	Drug Related Admission Rate (100%)	Alcohol Related Admission Rate (96%)	Alcohol Specific Death Rate (96%)
Belfast LGD	Drug Related Death Rate (96%)	Under 20 Teenage Birth Rate (91%)	Alcohol Related Admission Rate (90%)	Alcohol Specific Death Rate (83%)	Preventable Death Rate (80%)
Causeway Coast & Glens LGD	Alcohol Specific Death Rate (111%)	Drug Related Admission Rate (99%)	Smoking During Pregnancy (96%)	Alcohol Related Admission Rate (88%)	Under 20 Teenage Birth Rate (85%)
Derry City & Strabane LGD	Drug Related Death Rate (168%)	Alcohol Specific Death Rate (149%)	Alcohol Related Admission Rate (116%)	Drug Related Admission Rate (103%)	Lung Cancer Incidence Rate (92%)
Fermanagh & Omagh LGD	Self-Harm Admission Rate (67%)	Alcohol Related Admission Rate (64%)	Drug Related Admission Rate (61%)	Lung Cancer Death Rate (48%)	Lung Cancer Incidence Rate (46%)
Lisburn & Castlereagh LGD	Under 20 Teenage Birth Rate (155%)	Alcohol Specific Death Rate (139%)	Smoking During Pregnancy (101%)	Alcohol Related Admission Rate (95%)	Drug Related Admission Rate (92%)
Mid & East Antrim LGD	Drug Related Death Rate (220%)	Drug Related Admission Rate (157%)	Self-Harm Admission Rate (145%)	Smoking During Pregnancy (137%)	Alcohol Related Admission Rate (120%)
Mid Ulster LGD	Alcohol Related Admission Rate (59%)	Drug Related Admission Rate (43%)	Under 75 Death Rate – All Causes (26%)	Potential Years of Life Lost (24%)	Avoidable Death Rate (23%)
Newry, Mourne & Down LGD	Alcohol Specific Death Rate (68%)	Self-Harm Admission Rate (48%)	Alcohol Related Admission Rate (48%)	Drug Related Admission Rate (40%)	Preventable Mortality Rate (27%)

⁵⁶ Includes all indicators including outcomes relating to service-based indicators. For further information see 'Notes for users', Page 9

⁵⁷ Excludes gaps that were not statistically significant.

Antrim & Newtownabbey

Local Government District (LGD)

Comparison with NI⁵⁸

30 Health Outcomes were similar to the NI Average

2 Health Outcomes were better than the NI Average

Of the 32 health outcome indicators analysed for Antrim & Newtownabbey residents, none were worse than the NI average, 30 were similar to the NI average, and two of the health outcomes analysed were better than the NI average.

Life Expectancy:

In Antrim & Newtownabbey, male life expectancy (78.7 years) was similar to the NI average (78.4 years) and female life expectancy (82.5 years) was also similar to the average (82.3 years).

Gaps in Health Outcomes between Antrim & Newtownabbey LGD and NI⁵⁹

Life expectancy gaps at birth and at age 65 included in the figures above have not been displayed in the table below as they are represented in years.

Indicator	Gap	Comparison to NI
Primary 1 BMI: Obese	15%	Similar To
Primary 1 BMI: Overweight or Obese	11%	Similar To
Standardised Death Rate - COVID-19	8%	Similar To
Standardised Incidence Rate - Cancer	2%	Similar To
Standardised Death Rate - Alcohol Specific	2%	Similar To
Standardised Death Rate - Treatable	1%	Similar To
Standardised Prescription Rate - Mood & Anxiety	1%	Similar To
Standardised Death Rate - All Deaths	0%	Similar To
Standardised Death Rate - Cancer U75	0%	Similar To
Standardised Death Rate - Smoking Attributable Causes	-1%	Similar To
Standardised Death Rate - All Cause U75	-1%	Similar To
Standardised Incidence Rate - Lung Cancer	-1%	Similar To
Standardised Death Rate - Lung Cancer	-1%	Similar To
Standardised Prescription Rate - Antihypertensive	-2%	Better Than
Teenage Birth Rate U20	-2%	Similar To
Small for Gestational Age	-3%	Similar To
Standardised Death Rate - Avoidable	-3%	Similar To
Standardised Death Rate - Circulatory U75	-4%	Similar To
Standardised Death Rate - Preventable	-5%	Similar To
Year 8 BMI: Obese	-5%	Similar To
Breastfeeding on Discharge	-6%	Similar To
Potential Years of Life Lost	-6%	Similar To

⁵⁸ Outcomes relating to service-based indicators, including hospital admissions, have not been presented above as it is not possible to ascertain whether a LGD-NI gap in utilisation of services fully constitutes a difference in health outcomes due to potentially different service provision profiles across different geographic areas. For further information see 'Notes for users', Page 9.

⁵⁹ As confidence intervals vary across indicators, some gaps between the LGD and NI are not significant and have therefore been categorised as 'similar to', despite being larger than gaps for other indicators that have been identified as significant (i.e. worse than or better than).

Standardised Prescription Rate - Statin	-7%	Better Than
Year 8 BMI: Overweight or Obese	-8%	Similar To
Smoking During Pregnancy	-8%	Similar To
Low Birth Weight	-8%	Similar To
Standardised Death Rate - Drug Related Causes	-14%	Similar To
Crude Suicide Rate	-19%	Similar To

Comparison with Most Deprived Areas

Inequality gaps between the 20% most deprived areas within the Antrim & Newtownabbey LGD and the LGD average:

Life Expectancy

- Male life expectancy in the Trust's most deprived areas was 75.4 years, **3.3 years** less than the LGD average (78.7 years).
- Female life expectancy in the Trust's most deprived areas was 79.9 years, **2.7 years** less than the HSCT average (82.5 years).

Largest Inequality Gaps within Antrim & Newtownabbey LGD⁶⁰

Across the 51 indicators analysed, the majority of outcomes were significantly worse for those residing in the 20% most deprived areas of Antrim & Newtownabbey LGD when compared with the Antrim & Newtownabbey LGD average. Exceptions include under 75 circulatory death rate, deaths due to covid, suicide rate, under 20 teenage birth rate, low birth weight, babies born small for gestational age, and primary 1 & year 8 overweight and/or affected by obesity, where differences were not statistically significant. The largest inequality gaps observed in the latest year were⁶¹:

Drug Related Death Rate (114%)	Alcohol Specific Death Rate (111%)	Self-Harm Admission Rate (93%)	Smoking during Pregnancy (85%)	Drug Related Admission Rate (76%)
--------------------------------	------------------------------------	--------------------------------	--------------------------------	-----------------------------------

This is a summary of findings only. For a full assessment and all figures see downloadable tables at:

<https://www.health-ni.gov.uk/articles/health-inequalities-statistics>

⁶⁰ Includes all indicators including outcomes relating to service-based indicators. For further information see 'Notes for users', Page 9.

⁶¹ Excludes gaps that were not statistically significant.

Ards & North Down

Local Government District (LGD)

Comparison with NI⁶²

11 Health Outcomes were similar to the NI Average

21 Health Outcomes were better than the NI Average

Of the 32 health outcome indicators analysed for Ards & North Down residents, none were worse than the NI average, 11 were similar to the NI average, and 21 of the health outcomes analysed were better than the NI average.

Life Expectancy:

In Ards & North Down, male life expectancy (80.0 years) was 1.6 years higher than the NI average (78.4 years) and female life expectancy (82.9 years) was 0.6 years higher than the average (82.3 years).

Gaps in Health Outcomes between Ards & North Down LGD and NI⁶³

Life expectancy gaps at birth and at age 65 included in the figures above have not been displayed in the table below as they are represented in years.

Indicator	Gap	Comparison to NI
Low Birth Weight	10%	Similar To
Small for Gestational Age	3%	Similar To
Teenage Birth Rate U20	0%	Similar To
Year 8 BMI: Overweight or Obese	0%	Similar To
Standardised Prescription Rate - Antihypertensive	-1%	Similar To
Standardised Prescription Rate - Mood & Anxiety	-3%	Better Than
Standardised Incidence Rate - Cancer	-4%	Better Than
Standardised Death Rate - Drug Related Causes	-7%	Similar To
Standardised Death Rate - Cancer U75	-7%	Similar To
Standardised Death Rate - COVID-19	-8%	Similar To
Breastfeeding on Discharge	-8%	Similar To
Standardised Death Rate - All Deaths	-9%	Better Than
Standardised Prescription Rate - Statin	-9%	Better Than
Standardised Death Rate - Treatable	-10%	Better Than
Year 8 BMI: Obese	-10%	Similar To
Standardised Death Rate - All Cause U75	-11%	Better Than
Primary 1 BMI: Overweight or Obese	-12%	Better Than
Potential Years of Life Lost	-12%	Better Than
Standardised Death Rate - Lung Cancer	-13%	Better Than
Standardised Death Rate - Smoking Attributable Causes	-13%	Better Than
Standardised Death Rate - Circulatory U75	-14%	Similar To
Standardised Death Rate - Avoidable	-17%	Better Than

⁶² Outcomes relating to service-based indicators, including hospital admissions, have not been presented above as it is not possible to ascertain whether a LGD-NI gap in utilisation of services fully constitutes a difference in health outcomes due to potentially different service provision profiles across different geographic areas. For further information see 'Notes for users', Page 9.

⁶³ As confidence intervals vary across indicators, some gaps between the LGD and NI are not significant and have therefore been categorised as 'similar to', despite being larger than gaps for other indicators that have been identified as significant (i.e. worse than or better than).

Standardised Incidence Rate - Lung Cancer	-19%	Better Than
Standardised Death Rate - Preventable	-20%	Better Than
Smoking During Pregnancy	-21%	Better Than
Standardised Death Rate - Alcohol Specific	-25%	Better Than
Crude Suicide Rate	-26%	Better Than
Primary 1 BMI: Obese	-32%	Better Than

Comparison with Most Deprived Areas

Inequality gaps between the 20% most deprived areas within the Ards & North Down LGD and the LGD average:

Life Expectancy

- Male life expectancy in the Trust's most deprived areas was 77.2 years, **2.8 years** less than the LGD average (80.0 years).
- Female life expectancy in the Trust's most deprived areas was 81.0 years, **1.9 years** less than the HSCT average (82.9 years).

Largest Inequality Gaps within Ards & North Down LGD⁶⁴

Across the 51 indicators analysed, the majority of outcomes were significantly worse for those residing in the 20% most deprived areas of Ards & North Down LGD when compared with the Ards & North Down LGD average. Exceptions include female life expectancy at age 65, under 75 circulatory mortality, deaths due to covid, suicide rate, alcohol specific deaths, drug related deaths, under 20 teenage birth rate, breastfeeding on discharge, low birth weight, babies born small for gestational age, and primary 1 & year 8 overweight and/or affected by obesity, where differences were not statistically significant. The largest inequality gaps observed in the latest year were⁶⁵:

Drug Related Admission Rate (97%)	Smoking during Pregnancy (81%)	Alcohol Related Admission Rate (75%)	Preventable Death Rate (67%)	Self-Harm Admission Rate (64%)
-----------------------------------	--------------------------------	--------------------------------------	------------------------------	--------------------------------

This is a summary of findings only. For a full assessment and all figures see downloadable tables at: <https://www.health-ni.gov.uk/articles/health-inequalities-statistics>

⁶⁴ Includes all indicators including outcomes relating to service-based indicators. For further information see 'Notes for users', Page 9.

⁶⁵ Excludes gaps that were not statistically significant.

Armagh City, Banbridge & Craigavon

Local Government District (LGD)

Comparison with NI⁶⁶

20 Health Outcomes were similar to the NI Average

12 Health Outcomes were better than the NI Average

Of the 32 health outcome indicators analysed for Armagh City, Banbridge & Craigavon residents, none were worse than the NI average, 20 were similar to the NI average, and 12 of the health outcomes analysed were better than the NI average.

Life Expectancy:

In Armagh City, Banbridge & Craigavon, male life expectancy (78.8 years) was similar to the NI average (78.4 years) and female life expectancy (82.8 years) was 0.6 years higher than the average (82.3 years).

Gaps in Health Outcomes between Armagh City, Banbridge & Craigavon LGD and NI⁶⁷

Life expectancy gaps at birth and at age 65 included in the figures above have not been displayed in the table below as they are represented in years.

Indicator	Gap	Comparison to NI
Year 8 BMI: Obese	17%	Similar To
Year 8 BMI: Overweight or Obese	1%	Similar To
Standardised Prescription Rate - Statin	1%	Similar To
Standardised Prescription Rate - Antihypertensive	0%	Similar To
Primary 1 BMI: Overweight or Obese	-2%	Similar To
Low Birth Weight	-2%	Similar To
Breastfeeding on Discharge	-2%	Similar To
Standardised Death Rate - Treatable	-3%	Similar To
Standardised Death Rate - Circulatory U75	-3%	Similar To
Standardised Incidence Rate - Cancer	-4%	Better Than
Standardised Prescription Rate - Mood & Anxiety	-4%	Better Than
Potential Years of Life Lost	-4%	Similar To
Standardised Death Rate - All Deaths	-5%	Better Than
Standardised Incidence Rate - Lung Cancer	-6%	Similar To
Standardised Death Rate - Alcohol Specific	-7%	Similar To
Standardised Death Rate - Cancer U75	-7%	Similar To
Standardised Death Rate - Avoidable	-7%	Better Than
Standardised Death Rate - All Cause U75	-7%	Better Than
Crude Suicide Rate	-9%	Similar To
Standardised Death Rate - Preventable	-9%	Better Than
Standardised Death Rate - COVID-19	-9%	Similar To

⁶⁶ Outcomes relating to service-based indicators, including hospital admissions, have not been presented above as it is not possible to ascertain whether a LGD-NI gap in utilisation of services fully constitutes a difference in health outcomes due to potentially different service provision profiles across different geographic areas. For further information see 'Notes for users', Page 9.

⁶⁷ As confidence intervals vary across indicators, some gaps between the LGD and NI are not significant and have therefore been categorised as 'similar to', despite being larger than gaps for other indicators that have been identified as significant (i.e. worse than or better than).

Standardised Death Rate - Smoking Attributable Causes	-9%	Better Than
Standardised Death Rate - Lung Cancer	-10%	Better Than
Small for Gestational Age	-10%	Similar To
Smoking During Pregnancy	-14%	Similar To
Teenage Birth Rate U20	-15%	Similar To
Primary 1 BMI: Obese	-17%	Similar To
Standardised Death Rate - Drug Related Causes	-35%	Better Than

Comparison with Most Deprived Areas

Inequality gaps between the 20% most deprived areas within the Armagh City, Banbridge & Craigavon LGD and the LGD average:

Life Expectancy

- Male life expectancy in the Trust's most deprived areas was 75.9 years, **2.9 years** less than the LGD average (78.8 years).
- Female life expectancy in the Trust's most deprived areas was 81.6 years, **1.2 years** less than the HSCT average (82.8 years).

Largest Inequality Gaps within Armagh City, Banbridge & Craigavon LGD⁶⁸

Across the 51 indicators analysed, the majority of outcomes were significantly worse for those residing in the 20% most deprived areas of Armagh City, Banbridge & Craigavon LGD when compared with the Armagh City, Banbridge & Craigavon average. Exceptions include female life expectancy at birth and at age 65, under 75 cancer mortality, deaths due to covid, suicide rate, drug related mortality, breastfeeding on discharge, low birth weight, babies born small for gestational age, and primary 1 & year 8 overweight and/or affected by obesity, where differences were not statistically significant. The largest inequality gaps observed in the latest year were⁶⁹:

Under 20 Teenage Birth Rate (134%)	Smoking During Pregnancy (111%)	Drug Related Admission Rate (100%)	Alcohol Related Admission Rate (96%)	Alcohol Specific Death Rate (96%)
------------------------------------	---------------------------------	------------------------------------	--------------------------------------	-----------------------------------

This is a summary of findings only. For a full assessment and all figures see downloadable tables at:

<https://www.health-ni.gov.uk/articles/health-inequalities-statistics>

⁶⁸ Includes all indicators including outcomes relating to service-based indicators. For further information see 'Notes for users', Page 9.

⁶⁹ Excludes gaps that were not statistically significant.

Belfast

Local Government District (LGD)

Comparison with NI⁷⁰

24 Health Outcomes were worse than the NI Average

8 Health Outcomes were similar to the NI Average

Of the 32 health outcome indicators analysed for Belfast LGD residents, 24 were worse than the NI average, eight were similar to the NI average, and none of the health outcomes analysed were better than the NI average.

Life Expectancy:

In the Belfast LGD, male life expectancy (75.8 years) was 2.6 years less than the NI average (78.4 years) and female life expectancy (80.4 years) was 1.9 years less than the average (82.3 years).

Gaps in Health Outcomes between Belfast LGD and NI⁷¹

Life expectancy gaps at birth and at age 65 included in the figures above have not been displayed in the table below as they are represented in years.

Indicator	Gap	Comparison to NI
Standardised Death Rate - Drug Related Causes	89%	Worse Than
Teenage Birth Rate U20	68%	Worse Than
Smoking During Pregnancy	55%	Worse Than
Standardised Death Rate - Alcohol Specific	50%	Worse Than
Standardised Death Rate - Lung Cancer	48%	Worse Than
Crude Suicide Rate	47%	Worse Than
Standardised Death Rate - Preventable	43%	Worse Than
Standardised Incidence Rate - Lung Cancer	40%	Worse Than
Standardised Death Rate - Avoidable	38%	Worse Than
Potential Years of Life Lost	33%	Worse Than
Standardised Death Rate - All Cause U75	33%	Worse Than
Standardised Death Rate - Smoking Attributable Causes	30%	Worse Than
Standardised Death Rate - Cancer U75	28%	Worse Than
Standardised Death Rate - Treatable	27%	Worse Than
Standardised Death Rate - Circulatory U75	26%	Worse Than
Low Birth Weight	18%	Worse Than
Standardised Death Rate - All Deaths	16%	Worse Than
Standardised Prescription Rate - Mood & Anxiety	15%	Worse Than
Small for Gestational Age	14%	Similar To
Standardised Incidence Rate - Cancer	10%	Similar To
Standardised Prescription Rate - Antihypertensive	7%	Worse Than

⁷⁰ Outcomes relating to service-based indicators, including hospital admissions, have not been presented above as it is not possible to ascertain whether a LGD-NI gap in utilisation of services fully constitutes a difference in health outcomes due to potentially different service provision profiles across different geographic areas. For further information see 'Notes for users', Page 9.

⁷¹ As confidence intervals vary across indicators, some gaps between the LGD and NI are not significant and have therefore been categorised as 'similar to', despite being larger than gaps for other indicators that have been identified as significant (i.e. worse than or better than).

Primary 1 BMI: Obese	5%	Similar To
Standardised Death Rate - COVID-19	4%	Similar To
Breastfeeding on Discharge	4%	Similar To
Standardised Prescription Rate - Statin	3%	Worse Than
Year 8 BMI: Overweight or Obese	2%	Similar To
Primary 1 BMI: Overweight or Obese	0%	Similar To
Year 8 BMI: Obese	-1%	Similar To

Comparison with Most Deprived Areas

Inequality gaps between the 20% most deprived areas within the Belfast LGD and the LGD average:

Life Expectancy

- Male life expectancy in the Trust's most deprived areas was 70.8 years, **5.0 years** less than the LGD average (75.8 years).
- Female life expectancy in the Trust's most deprived areas was 76.9 years, **3.4 years** less than the HSCT average (80.4 years).

Largest Inequality Gaps within Belfast LGD⁷²

Across the 51 indicators analysed, the majority of outcomes were significantly worse for those residing in the 20% most deprived areas of Belfast LGD when compared with the Belfast LGD average.

Exceptions include deaths due to covid, babies born small for gestational age, and primary 1 & year 8 overweight and/or affected by obesity, where differences were not statistically significant. The largest inequality gaps observed in the latest year were⁷³:

Drug Related Death Rate (96%)	Under 20 Teenage Birth Rate (91%)	Alcohol Related Admission Rate (90%)	Alcohol Specific Death Rate (83%)	Preventable Death Rate (80%)
-------------------------------	-----------------------------------	--------------------------------------	-----------------------------------	------------------------------

This is a summary of findings only. For a full assessment and all figures see downloadable tables at:

<https://www.health-ni.gov.uk/articles/health-inequalities-statistics>

⁷² Includes all indicators including outcomes relating to service-based indicators. For further information see 'Notes for users', Page 9.

⁷³ Excludes gaps that were not statistically significant.

Causeway Coast & Glens

Local Government District (LGD)

Comparison with NI⁷⁴

1 Health Outcome was worse than the NI Average

21 Health Outcomes were similar to the NI Average

10 Health Outcomes were better than the NI Average

Of the 32 health outcome indicators analysed for Causeway Coast & Glens residents, one was worse than the NI average, 21 were similar to the NI average, and ten of the health outcomes analysed were better than the NI average.

Life Expectancy:

In Causeway Coast & Glens, male life expectancy (79.1 years) was 0.7 years higher than the NI average (78.4 years) and female life expectancy (82.6 years) was also similar to the average (82.3 years).

Gaps in Health Outcomes between Causeway Coast & Glens LGD and NI⁷⁵

Life expectancy gaps at birth and at age 65 included in the figures above have not been displayed in the table below as they are represented in years.

Indicator	Gap	Comparison to NI
Primary 1 BMI: Obese	31%	Similar To
Year 8 BMI: Obese	10%	Similar To
Primary 1 BMI: Overweight or Obese	10%	Similar To
Breastfeeding on Discharge	10%	Worse Than
Year 8 BMI: Overweight or Obese	9%	Similar To
Smoking During Pregnancy	9%	Similar To
Standardised Prescription Rate - Mood & Anxiety	0%	Similar To
Standardised Prescription Rate - Antihypertensive	-1%	Similar To
Standardised Death Rate - All Deaths	-3%	Similar To
Standardised Death Rate - Treatable	-3%	Similar To
Standardised Incidence Rate - Cancer	-4%	Better Than
Standardised Death Rate - Circulatory U75	-4%	Similar To
Standardised Prescription Rate - Statin	-4%	Better Than
Standardised Death Rate - Smoking Attributable Causes	-5%	Similar To
Low Birth Weight	-6%	Similar To
Standardised Death Rate - All Cause U75	-6%	Similar To
Teenage Birth Rate U20	-6%	Similar To
Potential Years of Life Lost	-7%	Similar To
Standardised Death Rate - Cancer U75	-7%	Similar To
Standardised Death Rate - Avoidable	-7%	Better Than

⁷⁴ Outcomes relating to service-based indicators, including hospital admissions, have not been presented above as it is not possible to ascertain whether a LGD-NI gap in utilisation of services fully constitutes a difference in health outcomes due to potentially different service provision profiles across different geographic areas. For further information see 'Notes for users', Page 9.

⁷⁵ As confidence intervals vary across indicators, some gaps between the LGD and NI are not significant and have therefore been categorised as 'similar to', despite being larger than gaps for other indicators that have been identified as significant (i.e. worse than or better than).

Standardised Death Rate - Preventable	-9%	Better Than
Standardised Incidence Rate - Lung Cancer	-11%	Better Than
Standardised Death Rate - Drug Related Causes	-12%	Similar To
Standardised Death Rate - COVID-19	-12%	Similar To
Standardised Death Rate - Lung Cancer	-12%	Better Than
Crude Suicide Rate	-19%	Similar To
Small for Gestational Age	-20%	Better Than
Standardised Death Rate - Alcohol Specific	-29%	Better Than

Comparison with Most Deprived Areas

Inequality gaps between the 20% most deprived areas within the Causeway Coast & Glens LGD and the LGD average:

Life Expectancy

- Male life expectancy in the Trust's most deprived areas was 76.3 years, **2.9 years** less than the LGD average (79.1 years).
- Female life expectancy in the Trust's most deprived areas was 80.3 years, **2.3 years** less than the HSCT average (82.6 years).

Largest Inequality Gaps within Causeway Coast & Glens LGD⁷⁶

Across the 51 indicators analysed, the majority of outcomes were significantly worse for those residing in the 20% most deprived areas of Causeway Coast & Glens LGD when compared with the Causeway Coast & Glens LGD average. Exceptions include male life expectancy at age 65, treatable mortality, under 75 cancer and circulatory death rates, cancer incidence rate, deaths due to covid, suicide rate, drug related deaths, low birth weight, babies born small for gestational age, and primary 1 & year 8 overweight and/or affected by obesity, where differences were not statistically significant. The largest inequality gaps observed in the latest year were⁷⁷:

Alcohol Specific Death Rate (111%)	Drug Related Admission Rate (99%)	Smoking During Pregnancy (96%)	Alcohol Related Admission Rate (88%)	Under 20 Teenage Birth Rate (85%)
------------------------------------	-----------------------------------	--------------------------------	--------------------------------------	-----------------------------------

This is a summary of findings only. For a full assessment and all figures see downloadable tables at:
<https://www.health-ni.gov.uk/articles/health-inequalities-statistics>

⁷⁶ Includes all indicators including outcomes relating to service-based indicators. For further information see 'Notes for users', Page 9.

⁷⁷ Excludes gaps that were not statistically significant.

Derry City & Strabane

Local Government District (LGD)

Comparison with NI⁷⁸

15 Health Outcomes were worse than the NI Average

17 Health Outcomes were similar to the NI Average

Of the 32 health outcome indicators analysed for Derry City & Strabane residents, 15 were worse than the NI average, 17 were similar to the NI average, and none of the health outcomes analysed were better than the NI average.

Life Expectancy:

In Derry City & Strabane, male life expectancy (77.5 years) was 0.9 years less than the NI average (78.4 years) and female life expectancy (81.5 years) was 0.7 years less than the average (82.3 years).

Gaps in Health Outcomes between Derry City & Strabane LGD and NI⁷⁹

Life expectancy gaps at birth and at age 65 included in the figures above have not been displayed in the table below as they are represented in years.

Indicator	Gap	Comparison to NI
Standardised Death Rate - Alcohol Specific	36%	Worse Than
Standardised Death Rate - COVID-19	26%	Similar To
Year 8 BMI: Obese	24%	Similar To
Standardised Incidence Rate - Lung Cancer	20%	Worse Than
Small for Gestational Age	19%	Similar To
Breastfeeding on Discharge	18%	Worse Than
Smoking During Pregnancy	18%	Similar To
Standardised Prescription Rate - Mood & Anxiety	18%	Worse Than
Standardised Death Rate - Smoking Attributable Causes	17%	Worse Than
Primary 1 BMI: Obese	16%	Similar To
Year 8 BMI: Overweight or Obese	14%	Similar To
Standardised Death Rate - Lung Cancer	13%	Worse Than
Standardised Prescription Rate - Statin	13%	Worse Than
Standardised Death Rate - Preventable	11%	Worse Than
Standardised Death Rate - Avoidable	10%	Worse Than
Standardised Death Rate - All Cause U75	8%	Worse Than
Standardised Death Rate - All Deaths	8%	Worse Than
Standardised Death Rate - Treatable	8%	Similar To
Potential Years of Life Lost	7%	Similar To
Low Birth Weight	7%	Similar To
Standardised Death Rate - Circulatory U75	7%	Similar To
Primary 1 BMI: Overweight or Obese	5%	Similar To

⁷⁸ Outcomes relating to service-based indicators, including hospital admissions, have not been presented above as it is not possible to ascertain whether a LGD-NI gap in utilisation of services fully constitutes a difference in health outcomes due to potentially different service provision profiles across different geographic areas. For further information see 'Notes for users', Page 9.

⁷⁹ As confidence intervals vary across indicators, some gaps between the LGD and NI are not significant and have therefore been categorised as 'similar to', despite being larger than gaps for other indicators that have been identified as significant (i.e. worse than or better than).

Standardised Incidence Rate - Cancer	4%	Worse Than
Standardised Death Rate - Cancer U75	3%	Similar To
Standardised Prescription Rate - Antihypertensive	1%	Similar To
Standardised Death Rate - Drug Related Causes	0%	Similar To
Crude Suicide Rate	0%	Similar To
Teenage Birth Rate U20	-14%	Similar To

Comparison with Most Deprived Areas

Inequality gaps between the 20% most deprived areas within the Derry City & Strabane LGD and the LGD average:

Life Expectancy

- Male life expectancy in the Trust's most deprived areas was 71.6 years, **6.0 years** less than the LGD average (77.5 years).
- Female life expectancy in the Trust's most deprived areas was 78.0 years, **3.5 years** less than the HSCT average (81.5 years).

Largest Inequality Gaps within Derry City & Strabane LGD⁸⁰

Across the 51 indicators analysed, the majority of outcomes were significantly worse for those residing in the 20% most deprived areas of Derry City & Strabane LGD when compared with the Derry City & Strabane LGD average. Exceptions include deaths due to covid, suicide rate, low birth weight, babies born small for gestational age, and primary 1 & year 8 overweight and/or affected by obesity, where differences were not statistically significant. The largest inequality gaps observed in the latest year were⁸¹:

Drug Related Death Rate (168%)	Alcohol Specific Death Rate (149%)	Alcohol Related Admission Rate (116%)	Drug Related Admission Rate (103%)	Lung Cancer Incidence Rate (92%)
--------------------------------	------------------------------------	---------------------------------------	------------------------------------	----------------------------------

This is a summary of findings only. For a full assessment and all figures see downloadable tables at:

<https://www.health-ni.gov.uk/articles/health-inequalities-statistics>

⁸⁰ Includes all indicators including outcomes relating to service-based indicators. For further information see 'Notes for users', Page 9.

⁸¹ Excludes gaps that were not statistically significant.

Fermanagh & Omagh

Local Government District (LGD)

Comparison with NI⁸²

1 Health Outcome was worse than the NI Average

23 Health Outcomes were similar to the NI Average

8 Health Outcomes were better than the NI Average

Of the 32 health outcome indicators analysed for Fermanagh & Omagh residents, one was worse than the NI average, 23 were similar to the NI average, and eight of the health outcomes analysed were better than the NI average.

Life Expectancy:

In Fermanagh & Omagh, male life expectancy (79.0 years) was similar to the NI average (78.4 years) and female life expectancy (82.8 years) was similar to the average (82.3 years).

Gaps in Health Outcomes between Fermanagh & Omagh LGD and NI⁸³

Life expectancy gaps at birth and at age 65 included in the figures above have not been displayed in the table below as they are represented in years.

Indicator	Gap	Comparison to NI
Crude Suicide Rate	18%	Similar To
Small for Gestational Age	13%	Similar To
Standardised Prescription Rate - Statin	13%	Worse Than
Standardised Death Rate - Alcohol Specific	7%	Similar To
Year 8 BMI: Overweight or Obese	5%	Similar To
Standardised Death Rate - Circulatory U75	0%	Similar To
Standardised Incidence Rate - Cancer	0%	Similar To
Standardised Incidence Rate - Lung Cancer	-2%	Similar To
Breastfeeding on Discharge	-2%	Similar To
Standardised Death Rate - Lung Cancer	-3%	Similar To
Standardised Prescription Rate - Antihypertensive	-4%	Better Than
Standardised Death Rate - COVID-19	-4%	Similar To
Standardised Death Rate - Smoking Attributable Causes	-4%	Similar To
Standardised Death Rate - All Deaths	-4%	Better Than
Potential Years of Life Lost	-5%	Similar To
Standardised Death Rate - Preventable	-5%	Similar To
Standardised Death Rate - Avoidable	-6%	Better Than
Primary 1 BMI: Overweight or Obese	-7%	Similar To
Standardised Death Rate - Cancer U75	-7%	Similar To
Standardised Death Rate - All Cause U75	-8%	Similar To
Standardised Death Rate - Treatable	-8%	Similar To

⁸² Outcomes relating to service-based indicators, including hospital admissions, have not been presented above as it is not possible to ascertain whether a LGD-NI gap in utilisation of services fully constitutes a difference in health outcomes due to potentially different service provision profiles across different geographic areas. For further information see 'Notes for users', Page 9.

⁸³ As confidence intervals vary across indicators, some gaps between the LGD and NI are not significant and have therefore been categorised as 'similar to', despite being larger than gaps for other indicators that have been identified as significant (i.e. worse than or better than).

Standardised Prescription Rate - Mood & Anxiety	-9%	Better Than
Primary 1 BMI: Obese	-13%	Similar To
Low Birth Weight	-17%	Better Than
Year 8 BMI: Obese	-18%	Similar To
Standardised Death Rate - Drug Related Causes	-27%	Similar To
Smoking During Pregnancy	-33%	Better Than
Teenage Birth Rate U20	-52%	Better Than

Comparison with Most Deprived Areas

Inequality gaps between the 20% most deprived areas within the Fermanagh & Omagh LGD and the LGD average:

Life Expectancy

- Male life expectancy in the Trust's most deprived areas was 76.5 years, **2.5 years** less than the LGD average (79.0 years).
- Female life expectancy in the Trust's most deprived areas was 80.5 years, **2.3 years** less than the HSC average (82.8 years).

Largest Inequality Gaps within Fermanagh & Omagh LGD⁸⁴

Across the 51 indicators analysed, the majority of outcomes were significantly worse for those residing in the 20% most deprived areas of Fermanagh & Omagh LGD when compared with the Fermanagh & Omagh LGD average. Exceptions include male life expectancy at age 65, treatable mortality, under 75 circulatory and cancer mortality, deaths due to covid, suicide rate, alcohol specific deaths, drug related deaths, smoking during pregnancy, under 20 teenage birth rate, breastfeeding on discharge, low birth weight, babies born small for gestational age, and primary 1 & year 8 overweight and/or affected by obesity, where differences were not statistically significant. The largest inequality gaps observed in the latest year were⁸⁵:

Self-Harm Admission Rate (67%)	Alcohol Related Admission Rate (64%)	Drug Related Admission Rate (61%)	Lung Cancer Death Rate (48%)	Lung Cancer Incidence Rate (46%)
--------------------------------	--------------------------------------	-----------------------------------	------------------------------	----------------------------------

This is a summary of findings only. For a full assessment and all figures see downloadable tables at:

<https://www.health-ni.gov.uk/articles/health-inequalities-statistics>

⁸⁴ Includes all indicators including outcomes relating to service-based indicators. For further information see 'Notes for users', Page 9.

⁸⁵ Excludes gaps that were not statistically significant.

Lisburn & Castlereagh

Local Government District (LGD)

Comparison with NI⁸⁶

10 Health Outcomes were similar to the NI Average

22 Health Outcomes were better than the NI Average

Of the 32 health outcome indicators analysed for Lisburn & Castlereagh residents, none were worse than the NI average, ten were similar to the NI average, and 22 of the health outcomes analysed were better than the NI average.

Life Expectancy:

In Lisburn & Castlereagh, male life expectancy (80.4 years) was 2.0 years higher than the NI average (78.4 years) and female life expectancy (83.1 years) was 0.8 years higher than the average (82.3 years).

Gaps in Health Outcomes between Lisburn & Castlereagh LGD and NI⁸⁷

Life expectancy gaps at birth and at age 65 included in the figures above have not been displayed in the table below as they are represented in years.

Indicator	Gap	Comparison to NI
Standardised Death Rate - COVID-19	11%	Similar To
Standardised Incidence Rate - Cancer	-5%	Better Than
Standardised Prescription Rate - Antihypertensive	-5%	Better Than
Small for Gestational Age	-5%	Similar To
Crude Suicide Rate	-7%	Similar To
Primary 1 BMI: Overweight or Obese	-7%	Similar To
Standardised Death Rate - Treatable	-7%	Similar To
Standardised Death Rate - All Deaths	-9%	Better Than
Low Birth Weight	-9%	Similar To
Standardised Prescription Rate - Statin	-9%	Better Than
Standardised Prescription Rate - Mood & Anxiety	-10%	Better Than
Standardised Death Rate - Cancer U75	-12%	Better Than
Standardised Death Rate - Smoking Attributable Causes	-16%	Better Than
Standardised Death Rate - Avoidable	-16%	Better Than
Teenage Birth Rate U20	-16%	Similar To
Year 8 BMI: Overweight or Obese	-17%	Better Than
Standardised Death Rate - All Cause U75	-17%	Better Than
Standardised Death Rate - Circulatory U75	-18%	Similar To
Potential Years of Life Lost	-20%	Better Than
Standardised Death Rate - Preventable	-20%	Better Than

⁸⁶ Outcomes relating to service-based indicators, including hospital admissions, have not been presented above as it is not possible to ascertain whether a LGD-NI gap in utilisation of services fully constitutes a difference in health outcomes due to potentially different service provision profiles across different geographic areas. For further information see 'Notes for users', Page 9.

⁸⁷ As confidence intervals vary across indicators, some gaps between the LGD and NI are not significant and have therefore been categorised as 'similar to', despite being larger than gaps for other indicators that have been identified as significant (i.e. worse than or better than).

Primary 1 BMI: Obese	-21%	Similar To
Breastfeeding on Discharge	-22%	Better Than
Standardised Death Rate - Lung Cancer	-25%	Better Than
Standardised Incidence Rate - Lung Cancer	-25%	Better Than
Standardised Death Rate - Alcohol Specific	-25%	Better Than
Standardised Death Rate - Drug Related Causes	-33%	Better Than
Smoking During Pregnancy	-38%	Better Than
Year 8 BMI: Obese	-43%	Better Than

Comparison with Most Deprived Areas

Inequality gaps between the 20% most deprived areas within the Lisburn & Castlereagh LGD and the LGD average:

Life Expectancy

- Male life expectancy in the Trust's most deprived areas was 77.2 years, **3.2 years** less than the LGD average (80.4 years).
- Female life expectancy in the Trust's most deprived areas was 78.9 years, **4.2 years** less than the HSCT average (83.1 years).

Largest Inequality Gaps within Lisburn & Castlereagh LGD⁸⁸

Across the 51 indicators analysed, the majority of outcomes were significantly worse for those residing in the 20% most deprived areas of Lisburn & Castlereagh LGD when compared with the Lisburn & Castlereagh average. Exceptions include under 75 circulatory mortality, deaths due to covid, suicide rate, drug related mortality, low birth weight, babies born small for gestational age, and primary 1 & year 8 overweight and/or affected by obesity, where differences were not statistically significant. The largest inequality gaps observed in the latest year were⁸⁹:

Under 20 Teenage Birth Rate (155%)	Alcohol Specific Death Rate (139%)	Smoking During Pregnancy (101%)	Alcohol Related Admission Rate (95%)	Drug Related Admission Rate (92%)
------------------------------------	------------------------------------	---------------------------------	--------------------------------------	-----------------------------------

This is a summary of findings only. For a full assessment and all figures see downloadable tables at:
<https://www.health-ni.gov.uk/articles/health-inequalities-statistics>

⁸⁸ Includes all indicators including outcomes relating to service-based indicators. For further information see 'Notes for users', Page 9.

⁸⁹ Excludes gaps that were not statistically significant.

Mid and East Antrim

Local Government District (LGD)

Comparison with NI⁹⁰

2 Health Outcomes were worse than the NI Average

27 Health Outcomes were similar to the NI Average

3 Health Outcomes were better than the NI Average

Of the 32 health outcome indicators analysed for Mid and East Antrim LGD residents, two were worse than the NI average, 27 were similar to the NI average, and three of the health outcomes analysed were better than the NI average.

Life Expectancy:

In the Mid and East Antrim LGD, male life expectancy (78.3 years) was similar to the NI average (78.4 years) and female life expectancy (82.5 years) was similar to the average (82.3 years).

Gaps in Health Outcomes between Mid and East Antrim LGD and NI⁹¹

Life expectancy gaps at birth and at age 65 included in the figures above have not been displayed in the table below as they are represented in years.

Indicator	Gap	Comparison to NI
Smoking During Pregnancy	34%	Worse Than
Primary 1 BMI: Obese	13%	Similar To
Teenage Birth Rate U20	6%	Similar To
Small for Gestational Age	6%	Similar To
Standardised Death Rate - Circulatory U75	6%	Similar To
Standardised Death Rate - Drug Related Causes	3%	Similar To
Low Birth Weight	2%	Similar To
Standardised Prescription Rate - Mood & Anxiety	2%	Worse Than
Year 8 BMI: Obese	2%	Similar To
Standardised Prescription Rate - Antihypertensive	1%	Similar To
Primary 1 BMI: Overweight or Obese	1%	Similar To
Standardised Death Rate - Treatable	0%	Similar To
Standardised Death Rate - All Deaths	0%	Similar To
Breastfeeding on Discharge	-1%	Similar To
Standardised Incidence Rate - Cancer	-2%	Similar To
Standardised Death Rate - All Cause U75	-3%	Similar To
Standardised Prescription Rate - Statin	-3%	Better Than
Potential Years of Life Lost	-3%	Similar To

⁹⁰ Outcomes relating to service-based indicators, including hospital admissions, have not been presented above as it is not possible to ascertain whether a LGD-NI gap in utilisation of services fully constitutes a difference in health outcomes due to potentially different service provision profiles across different geographic areas. For further information see 'Notes for users', Page 9.

⁹¹ As confidence intervals vary across indicators, some gaps between the LGD and NI are not significant and have therefore been categorised as 'similar to', despite being larger than gaps for other indicators that have been identified as significant (i.e. worse than or better than).

Year 8 BMI: Overweight or Obese	-4%	Similar To
Standardised Death Rate - Smoking Attributable Causes	-4%	Similar To
Standardised Death Rate - Avoidable	-5%	Similar To
Standardised Incidence Rate - Lung Cancer	-6%	Similar To
Standardised Death Rate - Preventable	-7%	Better Than
Standardised Death Rate - Cancer U75	-7%	Similar To
Standardised Death Rate - Alcohol Specific	-13%	Similar To
Standardised Death Rate - Lung Cancer	-15%	Better Than
Crude Suicide Rate	-19%	Similar To
Standardised Death Rate - COVID-19	-26%	Similar To

Comparison with Most Deprived Areas

Inequality gaps between the 20% most deprived areas within the Mid and East Antrim LGD and the LGD average:

Life Expectancy

- Male life expectancy in the Trust's most deprived areas was 71.7 years, **6.6 years** less than the LGD average (78.3 years).
- Female life expectancy in the Trust's most deprived areas was 78.7 years, **3.7 years** less than the HSCT average (82.5 years).

Largest Inequality Gaps within Mid and East Antrim LGD⁹²

Across the 51 indicators analysed, the majority of outcomes were significantly worse for those residing in the 20% most deprived areas of Mid and East Antrim LGD when compared with the Mid and East Antrim LGD average. Exceptions include under 75 cancer mortality, circulatory admissions, deaths due to covid, suicide rate, babies born small for gestational age, breastfeeding on discharge, and primary 1 & year 8 overweight and/or affected by obesity, where differences were not statistically significant. The largest inequality gaps observed in the latest year were⁹³:

Drug Related Death Rate (220%)	Drug Related Admission Rate (157%)	Self-Harm Admission Rate (145%)	Smoking During Pregnancy (137%)	Alcohol Related Admission Rate (120%)
--------------------------------	------------------------------------	---------------------------------	---------------------------------	---------------------------------------

This is a summary of findings only. For a full assessment and all figures see downloadable tables at:

<https://www.health-ni.gov.uk/articles/health-inequalities-statistics>

⁹² Includes all indicators including outcomes relating to service-based indicators. For further information see 'Notes for users', Page 9.

⁹³ Excludes gaps that were not statistically significant.

Mid Ulster

Local Government District (LGD)

Comparison with NI⁹⁴

1 Health Outcome was worse than the NI Average

14 Health Outcomes were similar to the NI Average

17 Health Outcomes were better than the NI Average

Of the 32 health outcome indicators analysed for Mid Ulster residents, one was worse than the NI average, 14 were similar to the NI average, and 17 of the health outcomes analysed were better than the NI average.

Life Expectancy:

In the Mid Ulster LGD, male life expectancy (78.9 years) was similar to the NI average (78.4 years) and female life expectancy (83.1 years) was 0.8 years higher than the average (82.3 years).

Gaps in Health Outcomes between Mid Ulster LGD and NI⁹⁵

Life expectancy gaps at birth and at age 65 included in the figures above have not been displayed in the table below as they are represented in years.

Indicator	Gap	Comparison to NI
Standardised Prescription Rate - Statin	7%	Worse Than
Breastfeeding on Discharge	1%	Similar To
Standardised Death Rate - Circulatory U75	0%	Similar To
Primary 1 BMI: Obese	-2%	Similar To
Standardised Death Rate - All Deaths	-3%	Similar To
Primary 1 BMI: Overweight or Obese	-3%	Similar To
Standardised Prescription Rate - Antihypertensive	-3%	Better Than
Small for Gestational Age	-6%	Similar To
Year 8 BMI: Obese	-6%	Similar To
Standardised Incidence Rate - Cancer	-6%	Better Than
Standardised Death Rate - Smoking Attributable Causes	-8%	Better Than
Standardised Death Rate - All Cause U75	-9%	Better Than
Potential Years of Life Lost	-10%	Better Than
Standardised Death Rate - Cancer U75	-10%	Better Than
Low Birth Weight	-10%	Better Than
Standardised Death Rate - Preventable	-11%	Better Than
Standardised Death Rate - Avoidable	-11%	Better Than
Standardised Death Rate - Treatable	-11%	Better Than
Standardised Prescription Rate - Mood & Anxiety	-12%	Better Than

⁹⁴ Outcomes relating to service-based indicators, including hospital admissions, have not been presented above as it is not possible to ascertain whether a LGD-NI gap in utilisation of services fully constitutes a difference in health outcomes due to potentially different service provision profiles across different geographic areas. For further information see 'Notes for users', Page 9.

⁹⁵ As confidence intervals vary across indicators, some gaps between the LGD and NI are not significant and have therefore been categorised as 'similar to', despite being larger than gaps for other indicators that have been identified as significant (i.e. worse than or better than).

Year 8 BMI: Overweight or Obese	-13%	Similar To
Standardised Death Rate - Lung Cancer	-16%	Better Than
Standardised Incidence Rate - Lung Cancer	-18%	Better Than
Crude Suicide Rate	-20%	Similar To
Smoking During Pregnancy	-22%	Better Than
Standardised Death Rate - COVID-19	-23%	Similar To
Standardised Death Rate - Alcohol Specific	-32%	Better Than
Teenage Birth Rate U20	-38%	Better Than
Standardised Death Rate - Drug Related Causes	-42%	Better Than

Comparison with Most Deprived Areas

Inequality gaps between the 20% most deprived areas within the Mid Ulster LGD and the LGD average:

Life Expectancy

- Male life expectancy in the Trust's most deprived areas was 77.9 years, **1.0 years** less than the LGD average (78.9 years).
- Female life expectancy in the Trust's most deprived areas was 81.5 years, **1.6 years** less than the HSC average (83.1 years).

Largest Inequality Gaps within Mid Ulster LGD⁹⁶

Across the 51 indicators analysed, the majority of outcomes were significantly worse for those residing in the 20% most deprived areas of Mid Ulster LGD when compared with the Mid Ulster LGD average. Exceptions include male & female life expectancies at birth and age 65, overall death rate, treatable mortality, under 75 cancer and circulatory death rates, circulatory admission rates, cancer incidence rate, deaths due to covid, elective inpatient admissions, self-harm admissions, suicide rate, alcohol specific mortality, smoking attributable mortality, lung cancer mortality, drug related mortality, smoking during pregnancy, under 20 teenage birth rate, breastfeeding on discharge rate, low birth weight, babies born small for gestational age, and primary 1 & year 8 overweight and/or affected by obesity, where differences were not statistically significant. The largest inequality gaps observed in the latest year were⁹⁷:

Alcohol Related Admission Rate (59%)	Drug Related Admission Rate (43%)	Under 75 Death Rate – All Causes (26%)	Potential Years of Life Lost (24%)	Avoidable Death Rate (23%)
--------------------------------------	-----------------------------------	--	------------------------------------	----------------------------

This is a summary of findings only. For a full assessment and all figures see downloadable tables at:

<https://www.health-ni.gov.uk/articles/health-inequalities-statistics>

⁹⁶ Includes all indicators including outcomes relating to service-based indicators. For further information see 'Notes for users', Page 9.

⁹⁷ Excludes gaps that were not statistically significant.

Newry, Mourne and Down

Local Government District (LGD)

Comparison with NI⁹⁸

1 Health Outcome was worse than the NI Average

24 Health Outcomes were similar to the NI Average

7 Health Outcomes were better than the NI Average

Of the 32 health outcome indicators analysed for Newry, Mourne and Down residents, one was worse than the NI average, 24 were similar to the NI average, and seven of the health outcomes analysed were better than the NI average.

Life Expectancy:

In Newry, Mourne and Down, male life expectancy (78.9 years) was similar to the NI average (78.4 years) and female life expectancy (82.6 years) was similar to the average (82.3 years).

Gaps in Health Outcomes between Newry, Mourne and Down LGD and NI⁹⁹

Life expectancy gaps at birth and at age 65 included in the figures above have not been displayed in the table below as they are represented in years.

Indicator	Gap	Comparison to NI
Standardised Death Rate - COVID-19 ¹⁰⁰	33%	Worse Than
Year 8 BMI: Overweight or Obese	6%	Similar To
Primary 1 BMI: Obese	6%	Similar To
Primary 1 BMI: Overweight or Obese	4%	Similar To
Standardised Death Rate - Cancer U75	2%	Similar To
Breastfeeding on Discharge	2%	Similar To
Year 8 BMI: Obese	2%	Similar To
Standardised Prescription Rate - Statin	0%	Similar To
Standardised Incidence Rate - Cancer	0%	Similar To
Standardised Prescription Rate - Antihypertensive	-1%	Similar To
Standardised Death Rate - All Deaths	-2%	Similar To
Crude Suicide Rate	-4%	Similar To
Standardised Prescription Rate - Mood & Anxiety	-4%	Better Than
Standardised Incidence Rate - Lung Cancer	-4%	Similar To
Potential Years of Life Lost	-5%	Similar To

⁹⁸ Outcomes relating to service-based indicators, including hospital admissions, have not been presented above as it is not possible to ascertain whether a LGD-NI gap in utilisation of services fully constitutes a difference in health outcomes due to potentially different service provision profiles across different geographic areas. For further information see 'Notes for users', Page 9.

⁹⁹ As confidence intervals vary across indicators, some gaps between the LGD and NI are not significant and have therefore been categorised as 'similar to', despite being larger than gaps for other indicators that have been identified as significant (i.e. worse than or better than).

¹⁰⁰ It is important to note that the rate of COVID deaths in NI in 2022 was much lower than in the previous two years and, while the rate in Newry, Mourne and Down LGD had a significantly higher rate than the NI average in 2022, the lower numbers present a greater chance of an inflated significant difference between areas. See page 94 for further information.

Standardised Death Rate - Lung Cancer	-6%	Similar To
Standardised Death Rate - Smoking Attributable Causes	-6%	Similar To
Low Birth Weight	-7%	Similar To
Standardised Death Rate - All Cause U75	-7%	Similar To
Standardised Death Rate - Preventable	-8%	Better Than
Standardised Death Rate - Avoidable	-10%	Better Than
Standardised Death Rate - Alcohol Specific	-12%	Similar To
Standardised Death Rate - Treatable	-14%	Better Than
Standardised Death Rate - Circulatory U75	-15%	Similar To
Small for Gestational Age	-16%	Similar To
Teenage Birth Rate U20	-16%	Similar To
Smoking During Pregnancy	-32%	Better Than
Standardised Death Rate - Drug Related Causes	-42%	Better Than

Comparison with Most Deprived Areas

Inequality gaps between the 20% most deprived areas within the Newry, Mourne and Down LGD and the LGD average:

Life Expectancy

- Male life expectancy in the Trust's most deprived areas was 76.4 years, **2.5 years** less than the LGD average (78.9 years).
- Female life expectancy in the Trust's most deprived areas was 82.5 years, **0.1 years** less than the HSC average (82.6 years).

Largest Inequality Gaps within Newry, Mourne and Down LGD¹⁰¹

Across the 51 indicators analysed, the majority of outcomes were significantly worse for those residing in the 20% most deprived areas of Newry, Mourne and Down LGD when compared with the Newry, Mourne and Down LGD average. Exceptions include female life expectancies at birth and age 65, treatable mortality, under 75 cancer and circulatory death rates, deaths due to covid, elective inpatient admission rate, suicide rate, smoking attributable mortality, drug related mortality, lung cancer incidence, smoking in pregnancy, under 20 teenage birth rate, breastfeeding on discharge, low birth weight, babies born small for gestational age, under 18 dental filling, extraction and registration rates, and primary 1 & year 8 overweight and/or affected by obesity, where differences were not statistically significant. The largest inequality gaps observed in the latest year were¹⁰²:

Alcohol Specific Death Rate (68%)	Self-Harm Admission Rate (48%)	Alcohol Related Admission Rate (48%)	Drug Related Admission Rate (40%)	Preventable Mortality Rate (27%)
-----------------------------------	--------------------------------	--------------------------------------	-----------------------------------	----------------------------------

This is a summary of findings only. For a full assessment and all figures see downloadable tables at:

<https://www.health-ni.gov.uk/articles/health-inequalities-statistics>

¹⁰¹ Includes all indicators including outcomes relating to service-based indicators. For further information see 'Notes for users', Page 9.

¹⁰² Excludes gaps that were not statistically significant.

APPENDICES

APPENDIX A: SOCIAL GRADIENT OF HEALTH

Health inequalities are often considered in terms of the gap between the most and least deprived quintiles of the population. However this does not account for those areas of intermediate levels of deprivation that may also be relatively disadvantaged in terms of their health status. The Marmot Review¹⁰³ demonstrated that there is a social gradient in health that runs from top to bottom of the socioeconomic spectrum, meaning that health inequalities affect everyone. There is consistent evidence from throughout the world that people at a socioeconomic disadvantage suffer a heavier burden of illness and have higher mortality rates than their better off counterparts.

Different inequality measures can give information about different aspects of inequalities. Some measures concentrate on the extremes of deprivation such as the most-least deprived (*or absolute*) gap analysis presented in the main body of this report, whilst others include relative inequality gaps across the socioeconomic scale – taking into account the whole population - and can give quite different interpretations of inequalities. Therefore, in addition to the most-least deprived (*or absolute*) gap analysis presented earlier in this report, a social gradient analysis using the Relative Index of Inequalities (RII) has been undertaken to provide a fuller assessment of inequalities.

Absolute gap (most-least deprived gap): This measure describes the absolute difference between the extremes of deprivation. It has the advantage that it is intuitive and straightforward to explain, but the disadvantage that, because it focuses only on the extremes of deprivation, it does not take account of patterns of inequalities observed across the intermediate groups.

Slope Index of Inequality (SII): SII describes the gradient of health observed across the deprivation scale. While the absolute gap shows the difference between two large groups, SII measures the difference in health outcomes between the theoretical most and least deprived individuals, according to linear regression across health outcomes for all deprivation deciles. SII therefore has the advantage of being sensitive to the experience of the entire population, rather than just the extremes of deprivation.

Relative Index of Inequality (RII): The RII describes the gradient of health observed across the deprivation scale, relative to the average for the observed population (by dividing the Slope of Index of Inequality (SII) by the mean). The value of RII tells you the magnitude of inequality in relation to the mean thus representing the proportionate change in the health outcome across the population. It allows inequalities to be compared and contrasted across a number of different health indicators, and also to be monitored over time.

For further information regarding the RII methodology, including how it is calculated, please refer to the NI Health & Social Care Inequalities Monitoring System – Regional 2014 report: <https://www.health-ni.gov.uk/publications/ni-health-and-social-care-inequalities-monitoring-system-hscims-regional-2014>

¹⁰³ Fair Society, Healthy Lives: The Marmot Review can be accessed at <http://www.instituteofhealthequity.org/resources-reports/fair-society-healthy-lives-the-marmot-review>


RESULTS

Social gradient analyses were carried out for the majority of indicators included in the HSCIMS. For some indicators this analysis could not be performed due to limitations on the level of data available. An explanatory interpretation of SII results is provided for life expectancy at birth below (it is not appropriate to look at relative gaps for life expectancy and therefore SII is used instead of RII), with a time series for the Absolute Gap (most-least deprived). It should be noted that in this report the SII value tends to be larger when compared with the absolute gap. This is due to the SII calculating across deprivation deciles whereas the absolute gap is calculated across deprivation quintiles, which can show different results. An explanatory interpretation of RII results is also provided for all cause mortality (U75), with RII then presented for all other indicators provided in Table 4.

Male Life Expectancy at Birth – SII

Year	2016-18	2017-19	2018-20	2019-21	2020-22
Absolute Gap (Most-Least Deprived)	7.1	7.0	6.9	7.3	7.2
Slope Index of Inequality (SII)	8.1	8.0	7.8	8.4	8.2

For life expectancy values, the Slope Index of Inequality (SII) indicates the absolute gap across the deprivation scale, represented by the gradient of the linear best fit line shown. In 2020-22, male life expectancy at birth indicates a SII gap of 8.2 years. This is higher than that indicated by the absolute gap between the most and least deprived quintiles (7.2 years). Both the absolute gap and SII showed no significant change in the male life expectancy deprivation gap between 2016-18 and 2020-22.



SDR – All Cause Mortality (U75) – RII

Year	2018	2019	2020	2021	2022
Absolute Gap (Most-Least Deprived)	114%	121%	116%	141%	117%
Slope Index of Inequality (SII)	-336	-345	-367	-404	-342
Relative Index of Inequality (RII)	-0.93	-0.98	-0.96	-1.06	-0.96

The Relative Index of Inequality (RII) indicates the relative gap across the deprivation scale. In 2022, mortality rates among those aged below 75 years indicated a deprivation gap of -0.96, meaning that the SII value of -342 deaths per 100,000 population is equivalent to 96% of the average under 75 mortality rate across NI. Both the RII and the absolute gap showed that the deprivation gap remained similar across the analysed period.




Table 4: Social Gradient Analysis of Indicators

A comparison of the inequality gaps provided in the main body of the report with social gradient results are presented in the table below:

Indicator		Time Series				
		2016-18	2017-19	2018-20	2019-21	2020-22
Male Life Expectancy at Birth						
Absolute Gap (Most–Least Deprived)	No Change	7.1	7.0	6.9	7.3	7.2
Slope Index of Inequality (SII)	No Change	8.1	8	7.8	8.4	8.2
Female Life Expectancy at Birth						
Absolute Gap (Most–Least Deprived)	No Change	4.4	4.9	5.0	5.1	4.8
Slope Index of Inequality (SII)	Widened	5.1	5.5	5.6	5.8	5.7
Male Life Expectancy at Age 65						
Absolute Gap (Most–Least Deprived)	Narrowed	3.2	3.1	3.1	3.0	2.9
Slope Index of Inequality (SII)	Narrowed	3.5	3.3	3.2	3.2	3.2
Female Life Expectancy at Age 65						
Absolute Gap (Most–Least Deprived)	Widened	2.4	2.6	2.7	2.7	2.8
Slope Index of Inequality (SII)	Widened	2.6	2.7	2.7	2.8	2.9
SDR-All Deaths						
Absolute Gap (Most–Least Deprived)	No Change	49%	45%	40%	52%	42%
Relative Index of Inequality (RII)	No Change	-0.46	-0.43	-0.38	-0.50	-0.43
Potential Years of Life Lost						
Absolute Gap (Most–Least Deprived)	No Change	126%	129%	125%	133%	130%
Relative Index of Inequality (RII)	Widened	-0.98	-1.01	-1.00	-1.05	-1.04
SDR- Treatable						
Absolute Gap (Most–Least Deprived)	No Change	89%	91%	86%	89%	83%
Relative Index of Inequality (RII)	No Change	-0.78	-0.78	-0.76	-0.77	-0.74
SDR – Preventable						
Absolute Gap (Most–Least Deprived)	Widened	182%	194%	191%	200%	202%
Relative Index of Inequality (RII)	Widened	-1.28	-1.31	-1.30	-1.32	-1.34
SDR – Avoidable						
Absolute Gap (Most–Least Deprived)	No Change	146%	155%	152%	159%	158%
Relative Index of Inequality (RII)	No Change	-1.11	-1.13	-1.12	-1.15	-1.15
SDR - Circulatory (U75)						
Absolute Gap (Most–Least Deprived)	No Change	134%	142%	136%	137%	127%
Relative Index of Inequality (RII)	No Change	-1.01	-1.04	-1.02	-1.04	-1.03
SDR - Respiratory (U75)						
Absolute Gap (Most–Least Deprived)	No Change	256%	248%	277%	252%	230%

Deprived)						
Relative Index of Inequality (RII)	No Change	-1.52	-1.51	-1.52	-1.51	-1.47
SDR - Cancer (U75)		2014-18	2015-19	2016-20	2017-21	2018-22
Absolute Gap (Most–Least Deprived)	No Change	69%	71%	70%	74%	74%
Relative Index of Inequality (RII)	No Change	-0.67	-0.66	-0.65	-0.66	-0.68
SDR - All Cause Mortality (U75)		2018	2019	2020	2021	2022
Absolute Gap (Most–Least Deprived)	No Change	114%	121%	116%	141%	117%
Relative Index of Inequality (RII)	No Change	-0.93	-0.98	-0.96	-1.06	-0.96
SAR - Circulatory		2016/17- 2018/19	2017/18- 2019/20	2018/19- 2020/21	2019/20- 2021/22	2020/21- 2022/23
Absolute Gap (Most–Least Deprived)	No Change	24%	22%	20%	19%	20%
Relative Index of Inequality (RII)	Narrowed	-0.24	-0.22	-0.21	-0.19	-0.17
SAR - Circulatory (U75)		2016/17- 2018/19	2017/18- 2019/20	2018/19- 2020/21	2019/20- 2021/22	2020/21- 2022/23
Absolute Gap (Most–Least Deprived)	No Change	35%	34%	34%	35%	36%
Relative Index of Inequality (RII)	No Change	-0.36	-0.35	-0.36	-0.35	-0.35
SPR - Antihypertensive		2018	2019	2020	2021	2022
Absolute Gap (Most–Least Deprived)	No Change	24%	21%	24%	24%	23%
Relative Index of Inequality (RII)	No Change	-0.25	-0.23	-0.26	-0.25	-0.25
SPR - Statin		2018	2019	2020	2021	2022
Absolute Gap (Most–Least Deprived)	No Change	33%	34%	35%	34%	32%
Relative Index of Inequality (RII)	No Change	-0.35	-0.35	-0.36	-0.35	-0.34
SAR - Respiratory		2016/17- 2018/19	2017/18- 2019/20	2018/19- 2020/21	2019/20- 2021/22	2020/21- 2022/23
Absolute Gap (Most–Least Deprived)	No Change	94%	95%	102%	103%	97%
Relative Index of Inequality (RII)	No Change	-0.75	-0.76	-0.81	-0.84	-0.81
SAR - Respiratory (U75)		2016/17- 2018/19	2017/18- 2019/20	2018/19- 2020/21	2019/20- 2021/22	2020/21- 2022/23
Absolute Gap (Most–Least Deprived)	No Change	113%	111%	121%	122%	119%
Relative Index of Inequality (RII)	Widened	-0.88	-0.88	-0.94	-0.97	-0.97
SIR - Cancer		2010-16	2011-17	2012-18	2013-19	2014-20
Absolute Gap (Most–Least Deprived)	No Change	22%	22%	22%	19%	21%
Relative Index of Inequality (RII)	No Change	-0.23	-0.23	-0.23	-0.21	-0.23
SDR - COVID-19				2020	2021	2022
Absolute Gap (Most–Least Deprived)				27%	90%	50%
Relative Index of Inequality (RII)				-0.25	-0.63	-0.62
SAR - All Admissions		2018/19	2019/20	2020/21	2021/22	2022/23
Absolute Gap (Most–Least Deprived)	No Change	40%	36%	39%	39%	35%
Relative Index of Inequality (RII)	No Change	-0.40	-0.37	-0.41	-0.40	-0.36
SAR - Emergency Admissions		2018/19	2019/20	2020/21	2021/22	2022/23

Absolute Gap (Most–Least Deprived)	Narrowed	64%	61%	62%	62%	52%
Relative Index of Inequality (RII)	Narrowed	-0.59	-0.56	-0.58	-0.58	-0.50
SAtR - Emergency Care Attendances		2018/19	2019/20	2020/21	2021/22	2022/23
Absolute Gap (Most–Least Deprived)	No Change	57%	55%	59%	57%	54%
Relative Index of Inequality (RII)	No Change	-0.52	-0.50	-0.54	-0.52	-0.51
SAR - Elective Inpatient Admissions		2018/19	2019/20	2020/21	2021/22	2022/23
Absolute Gap (Most–Least Deprived)	No Change	36%	28%	27%	21%	28%
Relative Index of Inequality (RII)	No Change	-0.36	-0.30	-0.31	-0.22	-0.30
SAR - Day Case Admissions		2018/19	2019/20	2020/21	2021/22	2022/23
Absolute Gap (Most–Least Deprived)	No Change	24%	19%	22%	26%	25%
Relative Index of Inequality (RII)	No Change	-0.26	-0.22	-0.28	-0.29	-0.26
SAR – Self-Harm Admissions		2014/15- 2018/19	2015/16- 2019/20	2016/17- 2020/21	2017/18- 2021/22	2018/19- 2022/23
Absolute Gap (Most–Least Deprived)	Narrowed	240%	216%	189%	170%	154%
Relative Index of Inequality (RII)	Narrowed	-1.51	-1.43	-1.31	-1.22	-1.17
Crude Suicide Rate		2016-18	2017-19	2018-20	2019-21	2020-22
Absolute Gap (Most–Least Deprived)	No Change	174%	128%	98%	115%	170%
Relative Index of Inequality (RII)	No Change	-1.26	-1.08	-0.94	-1.01	-1.18
SPR - Mood & Anxiety		2018	2019	2020	2021	2022
Absolute Gap (Most–Least Deprived)	No Change	68%	66%	67%	66%	66%
Relative Index of Inequality (RII)	No Change	-0.63	-0.63	-0.63	-0.62	-0.62
SAR - Alcohol Related Causes		2016/17- 2018/19	2017/18- 2019/20	2018/19- 2020/21	2019/20- 2021/22	2020/21- 2022/23
Absolute Gap (Most–Least Deprived)	Narrowed	300%	277%	276%	272%	250%
Relative Index of Inequality (RII)	Narrowed	-1.82	-1.75	-1.71	-1.68	-1.60
SDR - Alcohol Specific		2014-18	2015-19	2016-20	2017-21	2018-22
Absolute Gap (Most–Least Deprived)	No Change	315%	318%	306%	342%	334%
Relative Index of Inequality (RII)	Widened	-1.82	-1.84	-1.86	-1.89	-1.88
SDR - Smoking Attributable Causes		2014-18	2015-19	2016-20	2017-21	2018-22
Absolute Gap (Most–Least Deprived)	Widened	95%	97%	97%	99%	101%
Relative Index of Inequality (RII)	Widened	-0.79	-0.79	-0.79	-0.80	-0.82
SIR - Lung Cancer		2010-16	2011-17	2012-18	2013-19	2014-20
Absolute Gap (Most–Least Deprived)	No Change	150%	160%	163%	150%	155%
Relative Index of Inequality (RII)	No Change	-1.12	-1.16	-1.18	-1.16	-1.18
SDR - Lung Cancer		2014-18	2015-19	2016-20	2017-21	2018-22
Absolute Gap (Most–Least Deprived)	Narrowed	162%	160%	151%	152%	153%
Relative Index of Inequality (RII)	Narrowed	-1.22	-1.19	-1.17	-1.16	-1.17
SAR - Drug Related Causes		2016/17- 2018/19	2017/18- 2019/20	2018/19- 2020/21	2019/20- 2021/22	2020/21- 2022/23

Absolute Gap (Most–Least Deprived)	Narrowed	261%	240%	227%	216%	213%
Relative Index of Inequality (RII)	Narrowed	-1.60	-1.50	-1.46	-1.43	-1.47
SDR - Drug Related Causes		2014-18	2015-19	2016-20	2017-21	2018-22
Absolute Gap (Most–Least Deprived)	No Change	386%	399%	363%	307%	392%
Relative Index of Inequality (RII)	Narrowed	-2.12	-2.09	-2.02	-1.87	-1.98
SDR - Drug Misuse		2014-18	2015-19	2016-20	2017-21	2018-22
Absolute Gap (Most–Least Deprived)	Widened	425%	461%	412%	441%	494%
Relative Index of Inequality (RII)	No Change	-2.13	-2.14	-2.09	-2.16	-2.16
Smoking During Pregnancy		2018	2019	2020	2021	2022
Absolute Gap (Most–Least Deprived)	No Change	458%	340%	374%	440%	410%
Relative Index of Inequality (RII)	No Change	-1.77	-1.69	-1.72	-1.84	-1.90
Teenage Birth Rate (U20)		2018	2019	2020	2021	2022
Absolute Gap (Most–Least Deprived)	No Change	402%	514%	603%	360%	421%
Relative Index of Inequality (RII)	No Change	-1.85	-1.97	-2.02	-1.72	-1.69
Breastfeeding on Discharge		2018	2019	2020	2021	2022
Absolute Gap (Most–Least Deprived)	Narrowed	48%	46%	48%	47%	42%
Relative Index of Inequality (RII)	Narrowed	0.73	0.73	0.73	0.74	0.63
Low Birth Weight		2018	2019	2020	2021	2022
Absolute Gap (Most–Least Deprived)	Widened	38%	44%	50%	55%	54%
Relative Index of Inequality (RII)	No Change	-0.42	-0.53	-0.62	-0.63	-0.55
Small for Gestational Age			2019	2020	2021	2022
Absolute Gap (Most–Least Deprived)			78%	65%	65%	56%
Relative Index of Inequality (RII)			-0.74	-0.63	-0.66	-0.52
Primary 1 BMI: Obese		2018/19	2019/20	2020/21	2021/22	2022/23
Absolute Gap (Most–Least Deprived)		55%			129%	130%
Relative Index of Inequality (RII)		-0.46			-0.91	-0.76
Primary 1 BMI: Overweight or Obese		2018/19	2019/20	2020/21	2021/22	2022/23
Absolute Gap (Most–Least Deprived)		36%			48%	40%
Relative Index of Inequality (RII)		-1.25			-1.56	-1.54
Year 8 BMI: Obese		2018/19	2019/20	2020/21	2021/22	2022/23
Absolute Gap (Most–Least Deprived)		128%	127%			94%
Relative Index of Inequality (RII)		-0.46	-0.75			-0.76
Year 8 BMI: Overweight or Obese		2018/19	2019/20	2020/21	2021/22	2022/23
Absolute Gap (Most–Least Deprived)		54%	52%			47%
Relative Index of Inequality (RII)		-1.25	-2.01			-1.54
Standardised Dental Filling Rate - Total		2018/19	2019/20	2020/21	2021/22	2022/23

Absolute Gap (Most–Least Deprived)	No Change	48%	50%	22%	30%	44%
Relative Index of Inequality (RII)	No Change	-0.45	-0.46	-0.24	-0.31	-0.41
Standardised Dental Filling Rate - Total (U18)		2018/19	2019/20	2020/21	2021/22	2022/23
Absolute Gap (Most–Least Deprived)	No Change	64%	56%	27%	39%	57%
Relative Index of Inequality (RII)	No Change	-0.55	-0.49	-0.23	-0.35	-0.49
Standardised Dental Extraction Rate - Total		2018/19	2019/20	2020/21	2021/22	2022/23
Absolute Gap (Most–Least Deprived)	No Change	103%	105%	96%	109%	108%
Relative Index of Inequality (RII)	No Change	-0.82	-0.83	-0.78	-0.85	-0.84
Standardised Dental Extraction Rate - Total (U18)		2018/19	2019/20	2020/21	2021/22	2022/23
Absolute Gap (Most–Least Deprived)	Narrowed	16%	11%	1%	11%	7%
Relative Index of Inequality (RII)	Narrowed	-0.20	-0.10	0.05	-0.09	-0.04
Standardised Dental Crowning Rate - Total		2018/19	2019/20	2020/21	2021/22	2022/23
Absolute Gap (Most–Least Deprived)	No Change	54%	54%	8%	19%	35%
Relative Index of Inequality (RII)	No Change	-0.45	-0.48	-0.09	-0.20	-0.34
Standardised Dental Registration Rate		2018/19	2019/20	2020/21	2021/22	2022/23
Absolute Gap (Most–Least Deprived)	Narrowed	2%	3%	2%	1%	0%
Relative Index of Inequality (RII)	Narrowed	0.05	0.05	0.04	0.03	0.01
Standardised Dental Registration Rate (U18)		2018/19	2019/20	2020/21	2021/22	2022/23
Absolute Gap (Most–Least Deprived)	No Change	11%	11%	11%	11%	10%
Relative Index of Inequality (RII)	No Change	0.14	0.15	0.15	0.15	0.13

Changes in Inequality Gaps

In the majority of indicators, there was an agreement in the assessment of change between the absolute gap and the slope index of inequality or relative index of inequality.

The absolute deprivation gap of the following indicators remained constant while the social gradient analysis showed that the inequality widened:

Female Life Expectancy at birth

SDR - Alcohol Specific

Potential Years of Life Lost

SAR - Respiratory (U75)

The absolute deprivation gap of the following indicators remained constant while the social gradient analysis showed that the inequality narrowed:

SAR - Circulatory

SDR - Drug Related

The absolute deprivation gap of the following indicators widened while the social gradient analysis showed no change in the inequality gap:

Low Birth Weight

SDR - Drug Misuse

Ranking of Inequality Gaps

The table below displays, in rank order from largest to smallest, the eight indicators with the largest inequality gaps as identified by RII and absolute gap analysis. As can be seen, all eight indicators identified in each analysis were the same, with a few differences in the rank order of these inequality gaps. The 'Rank Change' column in the below table refers to how the ranking of indicators according to the size of the inequality gaps changes when analysed by RII compared with the absolute gap analysis. For example, in the below table SDR – Drug related Causes has the fourth largest absolute gap, but the second largest RII gap. Therefore, its rank change is 2 with the 'Increase' indicating the RII rank is increased (moved up 2 places) compared with the absolute gap rank.

Rank	Absolute Gap	RII	Rank Change
1	SDR - Drug Misuse	SDR - Drug Misuse	
2	Teenage Birth Rate (U20)	SDR - Drug Related Causes	2 Increase
3	Smoking During Pregnancy	Smoking During Pregnancy	
4	SDR - Drug Related Causes	SDR - Alcohol Specific	1 Increase
5	SDR - Alcohol Specific	Teenage Birth Rate (U20)	3 Decrease
6	SAR - Alcohol Related Causes	SAR - Alcohol Related Causes	
7	SDR - Respiratory (U75)	Primary 1 BMI: Overweight or Obese	
8	SAR - Drug Related Causes	SDR - Respiratory (U75)	1 Decrease

It should be noted that life expectancy gaps have not been included in the ranking of inequality gaps above. This is because proportionately, life expectancy gaps are comparatively lower to those ranking highest in the table above. However, as the gap refers to years of life, and as life expectancy is an overarching indicator of health status it is a vital statistic of high importance and reducing this gap is considered a high priority.¹⁰⁴


¹⁰⁴ Life expectancy inequality gaps are included as key overarching indicators of the public health strategic framework 'Making Life Better' www.health-ni.gov.uk/articles/making-life-better-strategic-framework-public-health

APPENDIX B: POPULATION ATTRIBUTABLE RISK (PAR) OF DEPRIVATION

Population Attributable Risk (PAR) measures the proportion of a disease/outcome (i.e. prevalence, mortality, admissions etc.) in the population that is attributable to deprivation and thus could be eliminated if deprivation were eliminated. This allows us to determine the proportional decrease in alcohol-related admissions in the population for example, in the hypothetical situation that all individuals had the same rate of alcohol-related admission as those in the highest socioeconomic category (least deprived deprivation decile). The PAR is calculated as the rate of disease in the overall population minus the rate in the unexposed group (least deprived).

PAR has been calculated in the table below for a number of health outcomes. As can be seen, the PAR percentage for preventable mortality in 2018-22 was 48% which indicates that almost half of preventable deaths in Northern Ireland were attributable to deprivation. This is reflected in the chart below showing the standardised death rate for preventable mortality by deprivation decile, with the proportion of deaths above the red line totalling the 48% of preventable deaths attributable to deprivation.

Indicator ¹⁰⁵	%PAR
SDR - Alcohol Related Causes (U75)	62%
Teenage Birth Rate (U20)	51%
SAR - Alcohol Related Causes	48%
SDR - Preventable	48%
SDR - Respiratory (U75)	42%
SDR - Avoidable	42%
SIR - Lung Cancer	41%
SAR - Self-Harm Admissions	41%
SDR - Circulatory (U75)	37%
CDR - Suicide	33%
SDR - Cancer (U75)	27%



¹⁰⁵ Caution should be taken as analysis includes indicators including outcomes relating to service-based indicators. For further information see 'Notes for users', Page 9

APPENDIX C: ADDITIONAL INDICATORS

The tables below refer to additional indicators which form part of the HSCIMS that have not been included in the main body of the report. For each indicator the figures are presented for NI, the 20% most deprived areas, the 20% least deprived areas and the most-least deprived inequality gap. In addition the RII is provided, where appropriate.

Median Fire Response Times ¹⁰⁶ Time (Minutes: Seconds)	2018/19	2019/20	2020/21	2021/22	2022/23
	All	All	All	All	All
NI	00:08:23	00:08:15	00:08:39	00:08:41	00:08:32
Most Deprived	00:06:40	00:06:42	00:07:00	00:06:57	00:06:44
Least Deprived	00:08:13	00:08:09	00:08:32	00:08:55	00:08:53
Most-Least Deprived	-19%	-18%	-18%	-22%	-24%

Median Ambulance Response Times ^{105,107,108,109} Time (Minutes: Seconds)	2019	2020	2021	2022	2023
	All	All	All	All	All
NI	00:15:3	00:17:1	00:24:3	00:23:1	00:25:1
Most Deprived	00:12:1	00:14:2	00:21:1	00:18:1	00:19:2
Least Deprived	00:16:2	00:17:5	00:27:3	00:24:3	00:27:3
Most-Least Deprived	-25%	-19%	-23%	-26%	-29%

Looked After Children Rate per 1,000 population under 18 years	2018	2019	2020	2021	2022
	All	All	All	All	All
NI	5.6	6.0	6.3	6.4	6.6
Most Deprived	11.2	10.7	12.3	13.5	14.2
Least Deprived	1.8	1.6	2.0	1.8	1.9
Most-Least Deprived	526%	575%	526%	648%	664%

¹⁰⁶ Evidence shows that emergency times are correlated more with location such as urban/rural than deprivation.

¹⁰⁷ In 2019/20, a new Clinical Response Model (CRM) programme was introduced along with a new set of ambulance categories in line with the national Ambulance Response Programme (ARP). Therefore, information from 2019/20 onwards is calculated differently and cannot be directly compared with previous years.

¹⁰⁸ COVID-19 surges may have influenced response times, through increased calls and/or greater staff absences rates.

¹⁰⁹ Please note that these statistics are presented for analysis of inequality gaps only and differ from the official statistics on Ambulance Response Times included within Department's annual '[Hospital Statistics: Emergency Care Activity](#)' report. Please refer to the official statistics when assessing response times in line with the Clinical Response Model targets.

Autism Prevalence in School Age Children ¹¹⁰	2018/19	2019/20	2020/21	2021/22	2022/23
Rate per 100,000 population	All	All	All	All	All
NI	3,331	4,237	4,490	4,675	5,039
10% Most Deprived	4,550	5,806	6,133	6,343	7,044
10% Least Deprived	3,200	3,986	4,101	4,012	4,191
10% Most-Least Deprived	42%	46%	50%	58%	68%
20% Most Deprived	4,321	5,515	5,890	6,185	6,920
20% Least Deprived	3,223	4,154	4,399	4,405	4,611
20% Most-Least Deprived	34%	33%	34%	40%	50%

Standardised Death Rate – All Cause (5-Year)	2014-18	2015-19	2016-20	2017-21	2018-22
Deaths per 100,000 population	All	All	All	All	All
NI	1,021	1,014	1,020	1,027	1,007
Most Deprived	1,251	1,257	1,261	1,276	1,257
Least Deprived	885	876	885	883	865
Most-Least Deprived	41%	44%	42%	45%	45%

Standardised Dental Filling Rate - Individuals	2018/19	2019/20	2020/21	2021/22	2022/23
Rate per 100,000 population	All	All	All	All	All
NI	21,389	20,446	6,951	11,706	15,610
Most Deprived	23,124	22,048	6,889	11,924	16,408
Least Deprived	19,301	18,256	6,391	10,579	13,868
Most-Least Deprived	20%	21%	8%	13%	18%

Standardised Dental Filling Rate - Individuals (U18)	2018/19	2019/20	2020/21	2021/22	2022/23
Rate per 100,000 population	All	All	All	All	All
NI	17,483	16,534	4,909	8,941	12,729
Most Deprived	19,677	18,569	4,976	9,252	14,039
Least Deprived	14,018	13,560	4,318	7,544	10,332
Most-Least Deprived	40%	37%	15%	23%	36%

Standardised Dental Extraction Rate - Individuals	2018/19	2019/20	2020/21	2021/22	2022/23
Rate per 100,000 population	All	All	All	All	All
NI	5,711	5,489	3,278	4,389	4,836
Most Deprived	7,604	7,389	4,390	5,988	6,558
Least Deprived	4,353	4,166	2,409	3,212	3,535
Most-Least Deprived	75%	77%	82%	86%	85%

¹¹⁰ Rates for the most and least deprived deciles and the 10% most and 10% Most-Least Deprived gap have also been provided as these are published within the Department of Health's official statistical report on Autism Prevalence in School Age Children 'The prevalence of autism (including Asperger's Syndrome) in school age children in Northern Ireland'.

Standardised Dental Extraction Rate - Individuals (U18)	2018/19	2019/20	2020/21	2021/22	2022/23
Rate per 100,000 population	All	All	All	All	All
NI	4,504	4,335	2,138	3,073	3,627
Most Deprived	4,940	4,684	2,163	3,325	3,819
Least Deprived	4,296	4,029	1,993	2,810	3,342
Most-Least Deprived	15%	16%	9%	18%	14%

Standardised Dental Crowning Rate - Individuals	2018/19	2019/20	2020/21	2021/22	2022/23
Rate per 100,000 population	All	All	All	All	All
NI	1,592	1,443	342	580	807
Most Deprived	1,923	1,792	355	633	956
Least Deprived	1,378	1,266	356	578	760
Most-Least Deprived	40%	42%	0%	9%	26%

APPENDIX D: URBAN-RURAL ANALYSIS

Urban-Rural analysis included below is based on the 2015 NISRA Urban-Rural classification, with the exception of Healthy Life Expectancy and Disability Free Life Expectancy which use the 2005 urban rural classification for all years up to and including 2015, due to data limitations. Further information regarding urban-rural classification can be found on the NISRA webpage at www.nisra.gov.uk/urban-rural-classification.

A positive inequality gap means that the health outcomes in urban areas are worse than in the rural areas.

Summary of findings

Compared with the regional average, urban areas experienced worse outcomes across the majority of indicators analysed, however fire and ambulance response times continue to remain higher in rural areas.

Outcomes that were significantly worse in urban areas than rural areas

Male Life Expectancy at Birth	SPR Antihypertensive	SDR Lung Cancer
Female Life Expectancy at Birth	SPR Statin	SAR Drug Related
Female Healthy Life Expectancy	SAR Respiratory	SDR Drug Related
Female Disability Free Life Expectancy	SAR Respiratory (U75)	SDR Drug Misuse
Male Disability Free Life Expectancy	SIR Cancer	Smoking During Pregnancy
Male Life Expectancy at 65	SAR All	Teenage Birth Rate
Female Life Expectancy at 65	SAR Emergency	Breastfeeding on Discharge
SDR – All Deaths	SaTr Emergency Care	Low Birth Weight
PYLL	SAR Day Case	Small for Gestational Age
SDR Treatable	SAR Self Harm	P1 Obese
SDR Preventable	CDR Suicide	P1 Overweight or Obese
SDR Avoidable	SPR Mood & Anxiety	Dental Fillings
SDR Circulatory (U75)	SAR Alcohol Related	Dental Extractions
SDR Respiratory (U75)	SDR Alcohol Specific	Dental Crownings
SDR Cancer (U75)	SDR Smoking Attributable	Dental Registrations
SDR All Cause (U75)	SIR Lung Cancer	Dental Registrations (U18)

Outcomes that were significantly better in urban areas than the rural areas

Ambulance Response Times	Dental Fillings (U18)	Dental Extractions (U18)
Fire Response Times		

Outcomes that were similar (or not significantly different) in urban and rural areas

Male Healthy Life Expectancy	SAR Circulatory (U75)	Y8 Obese
Infant Mortality Rate	SAR Elective Inpatient Admissions	Y8 Overweight or Obese
SAR Circulatory	SDR – COVID-19	

Figures for each indicator for NI, Rural areas, Urban areas, Mixed Urban-Rural areas and the NI-Rural Gap, are provided within the accompanying downloadable tables:

<https://www.health-ni.gov.uk/articles/health-inequalities-statistics>

APPENDIX E: TECHNICAL NOTES & DEFINITIONS

Indicators

There are 58 indicators included in the Northern Ireland analyses for the current report, with an additional 13 indicators included within Appendix C. Not all indicators are assessed at each level of geography (see Table 5), and dependent on the number of years data available, or any potential quality issues, assessments may not be made on all aspects of an indicator.

Due to random fluctuations in events over time, it is often necessary to aggregate more than one year of data for indicators, in order to ensure stability. The number of years of information that are required to aggregate for each indicator is informed by both the number of events and also an assessment of its annual variability.

Standardisation Methods

A number of indicators included in this report have been age standardised to allow the comparison of rates between populations with different age structures by relating them to a standard population, in this case the 2013 European Standard Population (90+ version). In most circumstances direct standardisation is used which not only allows the comparison of disease and death rates across both areas and time, but also to assess the relative burden of disease in a population.

Indicator Stability/Confidence Intervals

Indicator stability at the regional level does not mean that an indicator is also stable at the lower geographic levels of HSC Trust, LGD or DEA. To ensure robustness of the data, confidence intervals were calculated for rates for the most recent year at each geographic level, including the 20% most deprived Trust and LGD areas. The confidence interval for each standardised rate was assessed, in terms of its size and in relation to other comparable rates for other geographical areas, i.e. the Belfast Trust average and its 20% most deprived Trust areas. As a result of these assessments not all of the 62 indicators examined at the regional level were deemed robust enough to be presented at the sub-regional level, of these 58 indicators; 53 were found suitable to be published at the HSC Trust level, 51 at the LGD level and 44 at the DEA level. A full list of indicators and the level analysed can be found in [Table 5](#).

Confidence intervals are used to quantify the imprecision in the estimate of a particular value. Specifically it quantifies the imprecision that results from random variation in the estimation of the value. In many cases the source of this random variation is sampling, for example in Healthy Life Expectancy, as any measurement taken from a sample provides an imprecise estimate of the true population value. In public health many indicators are based on what can be considered to be complete data sets and not samples, e.g. age standardised mortality rates based on death registers. In these instances the imprecision arises not as a result of sampling variation but of 'natural' variation. The indicator is considered to be the outcome of a stochastic process, i.e. one which can be influenced by the random occurrences that are inherent in the world around us. In such instances the value actually observed is only one of the set that could occur under the same circumstances. Generally in public health, it is the underlying circumstances or process that is of interest and the actual value observed gives only an imprecise estimate of this 'underlying risk'.

Review of Suicide Statistics in Northern Ireland

Suicide deaths in Northern Ireland are defined as deaths from Self-inflicted Injury (also referred to as intentional self-harm) as well as Events of Undetermined Intent. This is consistent with the UK National Statistics definition. A death which is suspected to be suicide must be referred to the Coroner with the information provided by coroners at registration of the death used to code the underlying cause of death. In some instances, it can be difficult to establish whether the cause of death was suicide. If it is not clear, or the Coroner has not specifically stated that it is a suicide, these are coded as 'Undetermined'.

Following a quality exercise between NISRA Vital Statistics Unit and the Coroners' Service, to better understand drug related deaths and intent, improvements have been made in order to reduce the number of deaths coded as 'Undetermined'. This process highlighted that some deaths coded as 'Undetermined' would be better classified as 'Drug-related', 'Accidental' or 'Intentional self-harm and event of undetermined intent (Suicide)'. The review of suicide statistics was completed in autumn 2022. A statistical discontinuity now exists for these categories from 2015 onwards preventing comparisons with data that exists prior to the review. This limitation prevents complete sub-regional analysis over a five year period for the Suicide mortality indicator (CDR – Suicide) which uses five-year aggregated totals. Further information on this review and detailed statistics on the number of suicides registered each year in Northern Ireland can be accessed at the link below.

<https://www.nisra.gov.uk/publications/suicide-statistics>

Use of Suicide Statistics in this Publication

In this report, and other reports produced by PHIRB, mortality from suicide is calculated according to the UK National Statistics definition shown in the table below.

ICD-10 Underlying Cause Code	Description
X60-84, Y87.0	Self-inflicted Injury (Intentional self-harm)
Y10-Y34, Y87.2	Events of Undetermined Intent

Table 5: HSCIMS Indicators Analysed at Northern Ireland (NI), HSC Trust and LGD level

INDICATOR	NI	Trust	LGD
Male Life Expectancy at Birth	●	●	●
Female Life Expectancy at Birth	●	●	●
Male Life Expectancy at Age 65	●	●	●
Female Life Expectancy at Age 65	●	●	●
Male Healthy Life Expectancy	●		
Female Healthy Life Expectancy	●		
Male Disability Free Life Expectancy	●		
Female Disability Free Life Expectancy	●		
Standardised Death Rate – All Deaths	●	●	●
Potential Years of Life Lost –All	●	●	●
Standardised Death Rate – Treatable	●	●	●
Standardised Death Rate – Preventable	●	●	●
Standardised Death Rate – Avoidable	●	●	●
Standardised Death Rate - Circulatory (U75)	●	●	●
Standardised Death Rate - Respiratory (U75)	●	●	
Standardised Death Rate - Cancer (U75)	●	●	●
Standardised Death Rate - All Cause Mortality (U75)	●	●	●
Standardised Admission Rate –Circulatory	●	●	●
Standardised Admission Rate - Circulatory (U75)	●	●	●
Standardised Prescription Rate – Antihypertensive	●	●	●
Standardised Prescription Rate – Statin	●	●	●
Standardised Admission Rate – Respiratory	●	●	●
Standardised Admission Rate - Respiratory (U75)	●	●	●
Standardised Incidence Rate – Cancer	●	●	●
Standardised Death Rate – COVID-19	●	●	●
Standardised Admission Rate - All Admissions	●	●	●
Standardised Admission Rate - Emergency Admissions	●	●	●
Standardised Attendance Rate - Emergency Care	●	●	●
Standardised Admission Rate - Elective Inpatient Admissions	●	●	●
Standardised Admission Rate - Day Case Admissions	●	●	●
Standardised Admission Rate – Self-Harm Admissions	●	●	●
Crude Death Rate – Suicide	●	●	●
Standardised Prescription Rate - Mood & Anxiety	●	●	●
Standardised Admission Rate - Alcohol Related Causes	●	●	●
Standardised Death Rate - Alcohol Specific Causes	●	●	●
Standardised Death Rate - Smoking Attributable Causes	●	●	●
Standardised Incidence Rate - Lung Cancer	●	●	●
Standardised Death Rate - Lung Cancer	●	●	●
Standardised Admission Rate - Drug Related Causes	●	●	●
Standardised Death Rate - Drug Related Causes	●	●	●
Standardised Death Rate - Drug Misuse	●	●	
Infant Mortality Rate	●		
Smoking During Pregnancy	●	●	●
Teenage Birth Rate (U20)	●	●	●
Breastfeeding on Discharge	●	●	●
Low Birth Weight	●	●	●
Small for Gestational Age	●	●	●
Primary 1 BMI: Obese	●	●	●
Primary 1 BMI: Obese & Overweight	●	●	●
Year 8 BMI: Obese	●	●	●
Year 8 BMI: Obese & Overweight	●	●	●
Standardised Dental Filling Rate	●	●	●
Standardised Dental Filling Rate (U18)	●	●	●

INDICATOR	NI	Trust	LGD
Standardised Dental Extraction Rate	●	●	●
Standardised Dental Extraction Rate (U18)	●	●	●
Standardised Dental Crowning Rate	●	●	●
Standardised Dental Registration Rate	●	●	●
Standardised Dental Registration Rate (U18)	●	●	●

Methodology for Assessing Sub-regional Health Inequality Gaps

When making comparisons between Trust/LGD areas and the NI average, outcomes relating to service-based indicators, including hospital admissions, have been presented as 'higher than', 'lower than' or 'similar to'. For health outcome indicators, e.g., mortality rates, comparisons have been presented as 'better than', 'worse than' or 'similar to'. This is due to difficulties in ascertaining whether any differences in rates for service-based indicators are due to differences in demand (i.e., health of the population), or, because of access difficulties. All figures relating to service-based indicators should be treated with caution as they may also be impacted by external factors that are not reflective of service demand. As unmet demand is not accounted for in the data, these indicators should therefore be viewed as indicators of service provision rather than demand.

In order to provide an assessment of the Trust/LGD to NI inequality gaps, for the most recent year, analysis was performed to indicate whether the Trust/LGD average was better than, similar to, or worse than the NI average. Or in the case of service-based indicators, higher than, similar to or lower than.

If the Trust/LGD average of the health outcome had overlapping confidence intervals with the NI average, then the health outcome was reported as being similar to the NI average. Where confidence intervals did not overlap, the Trust/LGD average of the health outcome was reported as being either better or worse (or higher or lower) than the NI average. This methodology was employed for all standardised rates (i.e. death, admission, incidence, and prescription rates), life expectancies, and potential years of life lost..

For those health outcomes which did not have confidence intervals associated with them, such as teenage birth rate, a Z Score (range of +/- 2.5%) was calculated for each health outcome value and if this range overlapped when comparing two areas, the health outcomes were considered to be similar. This methodology was used to identify any health outcomes which were notably worse or better than the NI average. As with all of our observations of differences between areas and assessments of changes over time, conclusions are open to interpretation.

Mortality Rates

For simplicity of understanding, mortality figures are based on the single main underlying cause of death classification, but a death can be due to a variety of different causes. This can lead to an underestimation of the impact of common conditions associated with multiple causes of death (e.g. diabetes, influenza and pneumonia). All death figures used in the HSCIMS are based on the year that the death was registered and not necessarily the year in which the death occurred. While the vast majority of deaths are registered shortly after death, there may be a delay in registering some deaths. It is important to note that registration-based figures build in delays arising from system wide processes which can drive annual fluctuations in the series. Every death reported to the Coroner is carefully considered and is influenced by several factors specific to each case. These include whether the Coroner orders a post mortem, whether an inquest is required, the complexity of each case, and the number of cases reported to and being investigated by the Coroner at any point in time. Events such as infant death, drug related deaths or suicide are usually referred to a coroner and this legal process can take some time. Events such as infant death, drug related deaths or suicide are usually referred to a coroner and this legal process can take some time.

Population

Population is a vital part of rate calculations; a change to the size of a population or its age distribution will impact on rates and subsequently inequality gaps. For instance, overall yearly deaths in Northern Ireland remained between 14,903 and 17,159 from the turn of the century up to 2022, yet up to and including 2019, mortality rates had been generally falling – this can be partially explained by the growing and ageing Northern Ireland population. Between 2008 and 2022 for example, the population grew from 1,779,152 to 1,910,543; an increase of 131,391 persons (7.3%). During this time the proportion of the population aged 65 and over increased from 14% (247,500 persons) in 2008 to 19% (335,449 persons) in 2022.

Small Area Population Estimates


Population estimates at a relatively small geographic area level (i.e. Super Output Area (SOA) and Small Area (SA)), by age and gender, are used to calculate many of the HSCIMS indicators for deprivation quintiles and urban/rural. However, with the release of the 2021 NI Census, SAs and SOAs have been replaced by Data Zones (DZ) and Super Data Zones (SDZ) which do not map to the latest NI Multiple Deprivation Measure or Urban-Rural classification. As such, population estimates produced for NI are no longer available at this level and it has therefore been necessary for Information Analysis Directorate (IAD) to produce in-house SA and SOA estimates. This has been done by using the latest NISRA mid-year estimates for LGDs to inflate 2020 small area population estimates. These reworked estimates are validated by a process of integrity checks with higher level age and geography population totals published by NISRA.

Deprivation Classification

The deprivation classification used in this report is based on the Northern Ireland Multiple Deprivation Measure (NIMDM) produced by NISRA. The 20% most and least deprived areas are defined according to the NIMDM 2017.¹¹¹

Although the 2017 NIMDM is available at small area level it was decided to continue using the SOA classification within the HSCIMS to ensure continuity and comparability with the back series of data and across indicators. In addition, all analysis presented is based on multiple deprivation rather than any specific deprivation domain.

Chart 1 – 20% Most and least deprived areas in Northern Ireland according to the 2017 NIMDM



© Crown copyright and database rights NIMA MOU210

¹¹¹ <https://www.nisra.gov.uk/statistics/deprivation/northern-ireland-multiple-deprivation-measure-2017-nimdm2017>

Drug Related Admissions and Mortality

Please note that some observations may be due to changes in drug misuse behaviours among the population. There are ongoing concerns about polydrug misuse and the misuse of prescription drugs and new psychoactive substances. It appears that a significant cohort are engaging in increasingly risky behaviours, with an acute increase in related harms.

Childhood Obesity

The data cleansing parameters applied to the calculation of childhood obesity figures has been refined from 2017/18 onwards. Therefore figures should be treated with caution when making comparisons.

Year 8 BMI assessments have been reintroduced into this report after being temporarily removed for several years due to recording and data quality issues between 2018/19 and 2021/22. BMI assessments are not available for 2019/20 and 2020/21 school years for Primary 1 and, 2020/21 and 2021/22 school years for Year 8 BMI assessments, as the number of measurements recorded was negatively impacted by school closures due to the COVID-19 pandemic. Assessments of change have therefore not been provided. In addition, Year 8 BMI data for 2018/19 and 2019/20 does not include measurements across all HSC Trusts therefore figures for these years should be treated with caution.

Deaths due to COVID-19

The rate of deaths due to COVID-19 in NI in 2022 was much lower than in the previous two years. It is important to note this when assessing comparisons between different areas as the lower numbers present a greater chance of an inflated significant difference between areas. This was particularly evident the Newry, Mourne and Down LGD which had a significantly higher rate than the NI average in 2022.

To put this into context, the rate of deaths due to COVID-19 in 2022 in NI was 39 deaths per 100,000 population, and in Newry ,Mourne & Down LGD it was 52 deaths per 100,000 population (33% higher). However, in 2021 the rate in Newry ,Mourne & Down LGD was similar to the NI average and in 2020, the rate in Newry ,Mourne & Down LGD was 32% lower than the NI rate (all data included within the downloadable tables). This highlights the irregular pattern of this outcome at sub-regional levels which needs to be considered within the context of the latest figures.

Sources of Information

Table 6: Indicators and Supplementary Information

Information	Source
Deaths and births	General Register Office, Vital Statistics & Administrative Research and Support Branch (VARs), NI Statistics and Research Agency (NISRA)
Hospital Admissions	Hospital Inpatient System, Hospital Information Branch (HIB), Information Analysis Directorate (IAD), Department of Health (DoH)
Hospital Attendances	SYMPHONY & EEMS, Hospital Information Branch (HIB), Information Analysis Directorate (IAD), Department of Health (DoH)
General health & Longstanding limiting illness	NI Health Survey, Public Health Information & Research Branch (PHIRB), Information Analysis Directorate (IAD), Department of Health (DoH)
Prescriptions	Electronic Prescribing Eligibility System (EPES), Business Services Organisation (BSO)
Cancer Incidence	Northern Ireland Cancer Registry Live Database (collected from HSC Trusts), NI Cancer Registry (NICR)
Smoking during pregnancy, breastfeeding, low birth weight, small for gestational age, healthy birth weight	NI Maternity System (NIMATS)
Childhood overweight/obesity	Child Health System
Dental Indicators	Electronic Data Input (EDI), FPPS Dental Portal, HS45, Business Services Organisation (BSO)
Fire response times	NI Fire and Rescue Service (NIFRS) Data Hub, NIFRS
Ambulance response times	NI Ambulance Service (NIAS) Data Hub, NIAS
Interim Life Tables	Office for National Statistics (ONS)
NI Mid-Year Population Estimates	NI Statistics and Research Agency (NISRA)
NI Household Population Estimates	NI Statistics and Research Agency (NISRA)
European Standard Population (ESP) 2013	Eurostat
Deprivation classification	NI Multiple Deprivation Measure 2017 (NISRA)
Urban-rural classification	NI Statistics and Research Agency (NISRA)
Looked after Children	OC2 Community Information Return, Community Information Branch (CIB), Information Analysis Directorate (IAD), Department of Health (DoH)
Children with Autism	NI School Census, Community Information Branch (CIB), Information Analysis Directorate (IAD), Department of Health (DoH)

Indicator Definitions

Disease Classification - The indicators making up the HSCIMS are classified using the International Classification of Disease, 10th revision (ICD-10). This is the standard diagnostic tool for epidemiology, health management and clinical purposes, including the analysis of the general health situation of population groups.

A complete listing of ICD-10 codes can be found at the following web link:

www.who.int/classifications/apps/icd/icd10online/

LIFE EXPECTANCY	
Life Expectancy at Birth	The expected years of life at time of birth based on mortality patterns in the period in question. It is based on the average death rates over a three year period. Presented separately for males and females.
Life Expectancy at Age 65	The expected years of life at age 65 based on mortality patterns in the period in question. It is based on the average death rates over a three year period. Presented separately for males and females.
Healthy Life Expectancy (HLE)	<p>This is the average number of years a person can expect to live in good health. HLE provides an estimate of lifetime spent in 'Very Good' or 'Good' health, calculated using respondents' perception of their own health according to the Health Survey Northern Ireland (HSNI). HLE excludes communal establishments. All urban/rural analysis prior to 2016 is based on the 2005 urban-rural classification.</p> <p>Please note that due to the coronavirus (COVID-19) pandemic, data collection from the 2020/21 survey onwards moved from face-to-face interviewing to telephone mode. This may have influenced the responses given by respondents. In addition, the sample size has been lower as a result and children have not been included in the survey since 2019/20. As a result, data relating to children in 2019 has been held constant from 2020 onwards.</p>
Disability Free Life Expectancy (DFLE)	<p>This is the average number of years a person can expect to live disability free. DFLE provides an estimate of lifetime spent free from a limiting persistent (twelve months or more) illness or disability, based upon a self-rated functional assessment of health recorded in the HSNI. DFLE excludes communal establishments. All urban/rural analysis prior to 2016 is based on the 2005 urban-rural classification. It should be noted that the health survey question used to determine longstanding illness changed from 2012/13 onward by making specific reference to mental health conditions in addition to physical. The new question is based on the UK harmonised principle for long-lasting health conditions and illness. This change may have affected responses to the question and subsequently impacted on DFLE figures. For further information contact PHIRB (details on reverse of publication).</p> <p>Please note that due to the coronavirus (COVID-19) pandemic, data collection from the 2020/21 survey onwards moved from face-to-face interviewing to telephone mode. This may have influenced the responses given by respondents. In addition, the sample size has been lower as a result and children have not been included in the survey since 2019/20. As a result, data relating to children in 2019 has been held constant from 2020 onwards.</p>
Pregnancy & Early Years	
Teenage Birth Rate (U20)	The number of births in an area to teenage mothers (i.e. Between 13 and 19 years of age) expressed per 1,000 females.

Smoking during Pregnancy	The proportion of all live births, where the Health and Care Number (HCN) of the mother is recorded, that were to mothers that reported smoking during pregnancy. Data is based on smoking status as recorded at the earliest available antenatal booking appointment. As this indicator is self-reported, it may be subject to a degree of under-reporting. Figures for 2018 onwards have been revised due to a change in methodology and quality assurance processes, and therefore differ slightly from previous editions of this report.
Low Birth Weight	The proportion of all live births where the HCN of the mother is recorded and the birth weight of the child was less than 2,500g.
Breastfeeding on Discharge	The proportion of all live births, where the HCN of the mother is recorded, that were being breastfed on discharge from hospital. Figures include mothers' breastfeeding their child as well as using complementary feeding.
Small for Gestational Age	The proportion of all live births, where the HCN of the mother is recorded, that were small-for-gestational age (SGA). This is when an infant is born with a birth weight less than the 10th percentile, on a chart customised for maternal characteristics, for gestational age in body weight. Birth weight percentile is only available from 2019 onwards due to insufficient recording levels prior to 2019.

ADMISSIONS	
Hospital Information System (HIS)	Admissions data used to calculate rates are provided by the Hospital Information Branch and are extracted from the Hospital Information System (HIS). All mental health specialities have been excluded from the data. Figures are based on number of admissions and not individuals. Further information and definition on inpatient and day case activity is available at https://www.health-ni.gov.uk/articles/inpatient-and-day-case-activity .
Standardised Admission Rate (SAR)	This is calculated by standardising (using the direct method) the average admission rate in NI (over a predefined period) due to specified ICD-10 classification codes (may also be age specific) to the 2013 European Standard Population (ESP).
Indicator Name	
- All Admissions	Includes all acute inpatient and day case admissions (excluding regular day and night attenders, hospital transfers and other (maternity/delivery episodes)). Deaths and discharges have been used as an approximation for admissions.
- Emergency Admissions	A patient for whom admission is unpredictable and at short notice because of clinical need. All non-elective acute admissions excluding maternity, other and not known.
- Elective Inpatient Admissions	A patient for whom the decision to admit could be separated in time from the actual admission. Does not include day cases, not to be confused with elective admissions (which include day cases)
- Day Case Admissions	A patient admitted electively during the course of a day with the intention of receiving care who does not require the use of a hospital bed overnight and who returns home as scheduled. If this original intention is not fulfilled and the patient stays overnight, such a patient should be counted as an inpatient and is not counted as a day case admission.
- Circulatory	Selected according to International Classification of Disease (ICD-10) codes I00-I99 identified within the primary diagnostic field only. ¹¹²
- Circulatory U75	ICD-10 codes I00-I99, under 75 years of age (identified within the primary diagnostic field only).

¹¹² For a listing and explanation of topology or site codes see: International Statistical Classification of Diseases and Related Health Problems, Tenth Revision, World Health Organisation, Geneva. Or view online at: <http://apps.who.int/classifications/icd10/browse/2019/en#/I>

- Respiratory	ICD-10 codes J00-J99 (identified within the primary diagnostic field only).
- Respiratory U75	ICD-10 codes J00-J99, under 75 years of age (identified within the primary diagnostic field only).
- Alcohol Related Causes	Alcohol related causes included in Table 8 identified within the primary diagnostic field or any of the first 6 secondary diagnoses fields. Figures include inpatients and day cases and excludes regular attenders.
- Drug Related Causes	Drug related causes included in Table 10 identified within the primary diagnostic field or any of the first 6 secondary diagnoses fields. Figures include inpatients and day cases and excludes regular attenders.
- Self-Harm Admissions	ICD-10 codes X60-84 and Y87.0 identified within the primary diagnostic field or any of the first 6 secondary diagnoses fields. This indicator was developed to complement the suicide information, however it does not provide a complete picture of the problem of self-harm (or parasuicide) as in many instances, self-harm does not result in an acute admission to hospital. It should be noted that there have been a range of additional infrastructure provided to support people presenting with self-harm. These programmes may be contributing to the decrease in self-harm admissions.

ATTENDANCES

SYMPHONY & EEMS	Attendance data used to calculate rates are provided by the Hospital Information Branch and are from the two administrative systems used by emergency departments in Northern Ireland (SYMPHONY & EEMS). Figures are based on number of attendances and not individuals. Further information on emergency care activity is available at https://www.health-ni.gov.uk/articles/emergency-care-and-ambulance-statistics .
----------------------------	---

Indicator Name

- Emergency Care Attendances	New and unplanned review attendances at all Emergency Departments (Types 1 - 3). Data for RVH-RAES (Eye Casualty) not included prior to 2018/19. This relates to any patient who presents without appointment to an emergency care department. This differs from an emergency admission where a patient is admitted to an acute hospital by various routes, including through an emergency care department or via a General Practitioner. Rates for the Standardised Attendance Rate – Emergency Care have been revised in the current publication as a result of a quality assurance exercise. Values for previous years may therefore not match with earlier versions of the Health Inequalities Annual Report.
------------------------------	--

MORTALITY

Infant Mortality Rate	The number of infant deaths per 1,000 live births. Infant deaths refer to all deaths in the first year of life.
Potential Years of Life Lost (PYLL)	This is calculated by summing the deaths occurring at each age and multiplying this with the number of years a person of that age could have been expected to live. It is a summary measure of premature mortality, weighting deaths occurring at younger ages, which are, a priori, preventable. It uses the average age-specific life expectancy for each five year age band as the age to which a person in that age band might be expected to live.
Suicide (CDR) <i>Not age standardised as it was found to make little or no difference whilst introducing a new confidence interval</i>	The number of deaths by intentional self-harm and events of undetermined intent per 100,000 population ICD-10 codes X60-X84, Y87.0, Y10-Y34, Y87.2

Standardised Death Rate (SDR)	This is calculated by directly age standardising the average death rate in NI over a given period, due to specific causes of death (selected according to ICD-10 classification) to the 2013 European Standard Population (ESP). Some death rates relate to those under the age of 75 as indicators of premature mortality for specific diseases.
Indicator Name	
- All Cause U75	All causes, under 75 year of age
- All Age All Cause	All causes
- Treatable	Causes of death that can be mainly avoided through timely and effective health care interventions, including secondary prevention and treatment (i.e. after the onset of disease, to reduce case-fatality) – see Table 7 for full list of causes.
- Preventable	Causes of death that can be mainly avoided through effective public health and primary prevention interventions (i.e. before the onset of diseases/injuries, to reduce incidence) – see Table 7 for full list of causes.
- Avoidable	Avoidable deaths are all those defined as preventable and treatable – see Table 7 for full list of causes.
- Circulatory U75	ICD-10 codes I00-I99, under 75 year of age.
- Respiratory U75	ICD-10 codes J00-J99, under 75 year of age.
- Cancer U75	ICD-10 codes C00-C97, under 75 year of age.
- Lung Cancer	ICD-10 codes C33-C34.
- COVID-19	In this report deaths due to COVID-19 are defined as ‘deaths due to COVID-19’ and use the same International Classification of Disease Tenth Revision (ICD-10) codes as reported by NISRA within the Registrar General Annual Report i.e. ICD-10 codes U07.1, U07.2 and U10.9
- Alcohol Specific	Alcohol Specific causes – see Table 9 for full list of causes.
- Drug Related Causes	Drug related causes – see Table 10 for full list of causes.
- Drug Misuse	Deaths related to drug misuse – see Table 11 for full list of causes.
- Smoking Attributable Causes	Deaths due to Smoking attributable causes– see Table 12 for full list of causes.

Cancer Incidence

Northern Ireland Cancer Registry (NICR)	Cancer incidence numbers are extracted from the NICR's "live" database, and hence are continuously updated. As a result, an earlier extract taken at a later date may supply slightly different results. Therefore, although the overall trend will be the same, previously published data and data published elsewhere may have rates that vary slightly to what is published is here.
--	---

Northern Ireland Cancer Registry (NICR) Standardised Incidence Rate (SIR)	NICR publish official Standardised Incidence Rates (SIRs), however the HSCIMS publishes at further levels to allow for assessment of inequality gaps between different areas/population groups, including deprivation analysis. This is calculated by standardising (using the direct method) the average incidence rate in NI (over seven years) due to specified ICD-10 classification codes to the 2013 European Standard Population (ESP). As data is sourced from a live dataset that is subject to change, NICR annually provides the latest ten years of data to provide the latest picture. As a result, historical figures within this report are subject to change.
--	---

Indicator Name	ICD-10 Classification
- Cancer	ICD-10 codes C00-C97, excluding C44 (non-melanoma skin cancer which is quite common, in most cases easily treated and rarely fatal).
- Lung Cancer	ICD-10 codes C33 and C34.

Prescriptions	
Electronic Prescribing Eligibility System (EPES)	Prescription data is extracted from the EPES which is maintained by Business Services Organisation (BSO). The data provided covers drugs dispensed in primary care only, and includes prescriptions issued by all types of prescribers including doctors, nurses and dentists, and all those issued and dispensed by pharmacists, dispensing doctors and appliance suppliers. Drugs prescribed and dispensed in hospital cannot be captured centrally due to the use of different IT systems.
Standardised Prescription Rate (SPR)	This is calculated by standardising (using the direct method) the average prescription rate (over one year) in NI for people dispensed predefined prescription drugs, to the 2013 European Standard Population (ESP). Rates refer to number of persons prescribed a drug and does not include multiple prescription.
Indicator Name	British National Formulary (BNF) code
- Antihypertensive	Drugs included are those with a BNF code of 2.2.1, 2.4, 2.5.5.1, 2.5.5.2 and 2.6.2
- Statin	Drugs included are those with a BNF code of 2.12
- Mood & Anxiety Disorders	Drugs included are those with a BNF code of 4.1.2 and 4.3
Diet and Dental Health	
Childhood Overweight and Obese	Height and weight information is extracted from the Child Health System (CHS) and converted into a Body Mass Index (BMI) score for each pupil. The BMI can be categorized using International Growth Charts as determined by the International Obesity Taskforce (IOTF) which consider age and gender, allowing the identification of those who are overweight or obese. Records are analysed based on two criteria: Date of Exam within the Primary 1 or Year 8 school year: 01/09/XX-31/08/XX Date of Birth for Primary 1 or Year 8 pupils: 02/07/XX - 01/07/XX For data since 2017/18, additional data cleansing parameters have been applied.
Dental indicators	Data on the number of fillings, extractions, crownings and individuals registered with a dentist are supplied by Business Services Organisation (BSO). Dental indicators are age standardised using population data sourced from NISRA, as with the majority of other indicators used in this publication. Dental indicators for fillings, extractions and crownings only includes paid treatments carried out by General Dental Service (GDS) dentists. These indicators do not include private work or secondary care activity, including work carried out by the Community Dental Service. Please note that for each dental indicator there is a possibility that individuals may be counted in two age groups, if they had a birthday between two or more separate procedures within the same year. This occurs in only a small number of cases (typically around 2% or less), but should be considered when interpreting data.
Standardised dental indicator rate	This is calculated by standardising (using the direct method) the indicator value in NI (over a predefined period) to the 2013 ESP.
Indicator Name	CHS Data
- Primary 1 BMI: Obese	The proportion of children in Primary 1 classified as obese.
- Primary 1 BMI: Overweight or Obese	The proportion of children in Primary 1 classified as overweight or obese.

- Year 8 BMI: Obese	The proportion of children in Year 8 classified as obese.
- Year 8 BMI: Overweight or Obese	The proportion of children in Year 8 classified as overweight or obese.
Indicator Name	BSO Data
- Standardised Dental Filling Rate	Total number of fillings per 100,000 population.
- Standardised Dental Filling Rate (U18)	Total number of fillings in under 18s per 100,000 population.
- Standardised Dental Filling Rate – Individuals	Individuals receiving one or more fillings per 100,000 population.
- Standardised Dental Filling Rate – Individuals (U18)	Individuals aged under 18 receiving one or more fillings per 100,000 population.
- Standardised Dental Extraction Rate	Total number of extractions per 100,000 population.
- Standardised Dental Extraction Rate (U18)	Total number of extractions in under 18s per 100,000 population.
- Standardised Dental Extraction Rate – Individuals	Individuals receiving one or more extractions per 100,000 population.
- Standardised Dental Extraction Rate – Individuals (U18)	Individuals aged under 18 receiving one or more extractions per 100,000 population.
- Standardised Dental Crowning Rate	Total number of crownings per 100,000 population.
- Standardised Dental Crowning Rate – Individuals	Individuals receiving one or more crownings per 100,000 population.
- Standardised Dental Registration Rate	Individuals registered with a dentist per 100,000 population
- Standardised Dental Registration Rate (U18)	Individuals aged under 18 registered with a dentist per 100,000 population

Additional Indicators	
Median Fire Response Time	<p>The median response time taken by the Northern Ireland Fire and Rescue Service (NIFRS) to respond to an incident. The 'response time' is measured as the 'time of the call to NIFRS Regional Control Centre' to 'the time the 1st Appliance books in attendance' at the incident.</p> <p>Calculations are based on the time taken for NIFRS to respond to each incident within a one year time period.</p> <p>The median i.e. midpoint value is reported rather than the simple average as it is unaffected by atypically long or short response times.</p> <p>For consistency the methodology used in this report is the one used in previous years. However the measuring of response times has been revised for internal reports. When measuring performance internally the Data Hub removes any incidents where the response time is less than one minute or greater than one hour to avoid outliers skewing performance.</p>
Median Ambulance Response Time	<p>The median time taken by the appropriate response vehicle to respond to an incident. Calculations are based on the time taken to respond to each incident within a one month time period (August). This data refers to CAT1, CAT1(T) and CAT2 emergency responses, excluding Healthcare Professionals (HCP) calls.</p> <p>The median i.e. midpoint value is reported rather than the simple average as it is unaffected by atypically long or short response times.</p>

Looked after Children	The number of looked after children per 100,000 population (under 18 years of age) by location prior to last entering care. Data was extracted from the annual OC2 Community Information Return, which includes children who have been in care continuously for twelve months or longer at 30th September.
Autism Prevalence in School Age Children	The number of children with Autism or Asperger Syndrome per 100,000 children in compulsory grant-aided education. Data extracted from the NI School Census.

ICD-10 Classification Tables

The table below lists the revised ICD-10 classification codes of all causes of death considered avoidable, with indication as to which are considered treatable, preventable or both. This definition was implemented following an Office for National Statistics (ONS) consultation,¹¹³ on the latest definition of avoidable mortality as proposed by the Organisation for Economic Co-operation and Development (OECD).¹¹⁴ All avoidable mortality indicators, including the back series, are based on the new definition. The previous definition, for use with the additional indicators in [Appendix C](#), can be found in Table 13. Within the OECD definition, recently COVID-19 has been added to the definition in the new category “Provisional assignment of new diseases”.

Table 7: Treatable, Preventable & Avoidable Causes

Condition Group & Cause	ICD-10 Codes	Age	Treatable	Preventable
Infectious Diseases				
Intestinal Diseases	A00-A09	0-74		•
Diphtheria, Tetanus, Poliomyelitis	A35, A36, A80	0-74		•
Whooping cough	A37	0-74		•
Meningococcal infection	A39	0-74		•
Sepsis due to streptococcus pneumonia and sepsis due to haemophilus influenza	A40.3, A41.3	0-74		•
Haemophilus influenza infections	A49.2	0-74		•
Sexually transmitted infections (except HIV/AIDS)	A50-A60, A63, A64	0-74		•
Varicella	B01	0-74		•
Measles	B05	0-74		•
Rubella	B06	0-74		•
Viral Hepatitis	B15-B19	0-74		•
HIV/AIDS	B20-B24	0-74		•
Malaria	B50-B54	0-74		•
Haemophilus and pneumococcal meningitis	G00.0, G00.1	0-74		•
Tuberculosis	A15-A19, B90, J65	0-74	• (50%)	• (50%)
Scarlet fever	A38	0-74	•	
Sepsis	A40 (excl. A40.3), A41 (excl. A41.3)	0-74	•	
Cellulitis	A46, L03	0-74	•	
Legionnaires disease	A48.1	0-74	•	
Streptococcal and enterococci infection	A49.1	0-74	•	
Other meningitis	G00.2, G00.3, G00.8, G00.9	0-74	•	
Meningitis due to other and unspecified causes	G03	0-74	•	

¹¹³ <https://www.ons.gov.uk/peoplepopulationandcommunity/healthandsocialcare/causesofdeath/methodologies/avoidablemortalityinenglandandwalesqmi#important-points>

¹¹⁴ <http://www.oecd.org/health/health-systems/Avoidable-mortality-2019-Joint-OECD-Eurostat-List-preventable-treatable-causes-of-death.pdf>

Condition Group & Cause	ICD-10 Codes	Age	Treatable	Preventable
Cancer				
Lip, oral cavity and pharynx cancer	C00-C14	0-74		•
Oesophageal cancer	C15	0-74		•
Stomach cancer	C16	0-74		•
Liver cancer	C22	0-74		•
Lung cancer	C33-C34	0-74		•
Mesothelioma	C45	0-74		•
Skin (melanoma) cancer	C43	0-74		•
Bladder cancer	C67	0-74		•
Cervical cancer	C53	0-74	• (50%)	• (50%)
Colorectal cancer	C18-C21	0-74	•	
Breast cancer (Female only)	C50	0-74	•	
Uterus cancer	C54,C55	0-74	•	
Testicular cancer	C62	0-74	•	
Thyroid cancer	C73	0-74	•	
Hodgkin's disease	C81	0-74	•	
Lymphoid leukaemia	C91.0, C91.1	0-74	•	
Benign neoplasm	D10-D36	0-74	•	
Endocrine and metabolic diseases				
Nutritional deficiency anaemia	D50-D53			•
Diabetes mellitus	E10-E14		• (50%)	• (50%)
Thyroid disorders	E00-E07		•	
Adrenal disorders	E24-E25 (except E24.4), E27		•	
Diseases of the nervous system				
Epilepsy	G40,G41	0-74	•	
Diseases of the circulatory system				
Aortic aneurysm	I71	0-74	• (50%)	• (50%)
Hypertensive diseases	I10-I13, I15	0-74	• (50%)	• (50%)
Ischaemic heart diseases	I20-I25	0-74	• (50%)	• (50%)
Cerebrovascular diseases	I60-I69	0-74	• (50%)	• (50%)
Other atherosclerosis	I70, I73.9	0-74	• (50%)	• (50%)
Rheumatic and other heart diseases	I00-I09	0-74	•	
Venous thromboembolism	I26, I80, I82.9	0-74	•	

Condition Group & Cause	ICD-10 Codes	Age	Treatable	Preventable
Diseases of the respiratory system				
Influenza	J09-J11	0-74		•
Pneumonia due to Streptococcus pneumoniae or Haemophilus influenza	J13-J14	0-74		•
Chronic lower respiratory diseases	J40-J44	0-74		•
Lung diseases due to external agents	J60-J64, J66-J70, J82, J92	0-74		•
Upper respiratory infections	J00-J06, J30-J39	0-74	•	
Pneumonia, not elsewhere classified or organism unspecified	J12, J15, J16- J18	0-74	•	
Acute lower respiratory infections	J20-J22	0-74	•	
Asthma and bronchiectasis	J45-J47	0-74	•	
Adult respiratory distress syndrome	J80	0-74	•	
Pulmonary oedema	J81	0-74	•	
Abscess of lung and mediastinum pyothorax	J85, J86	0-74	•	
Other pleural disorders	J90, J93, J94	0-74	•	
Diseases of the digestive system				
Gastric and duodenal ulcer	K25-K28	0-74	•	
Appendicitis	K35-K38	0-74	•	
Abdominal hernia	K40-K46	0-74	•	
Cholelithiasis and cholecystitis	K80-K81	0-74	•	
Other diseases of gallbladder or biliary tract	K82-K83	0-74	•	
Acute pancreatitis	K85.0, K85.1, K85.3, K85.8, K85.9	0-74	•	
Other diseases of pancreas	K86.1, K86.2, K86.3, K86.8, K86.9	0-74	•	
Diseases of the genitourinary system				
Nephritis and nephrosis	N00-N07	0-74	•	
Obstructive uropathy	N13, N20-N21, N35	0-74	•	
Renal failure	N17-N19	0-74	•	
Renal colic	N23	0-74	•	
Disorders resulting from renal tubular dysfunction	N25	0-74	•	
Unspecified contracted kidney, small kidney of unknown cause	N26-N27	0-74	•	
Inflammatory diseases of genitourinary system	N34.1, N70-N73, N75.0, N75.1, N76.4, N76.6	0-74	•	
Prostatic hyperplasia	N40	0-74	•	

Condition Group & Cause	ICD-10 Codes	Age	Treatable	Preventable
Pregnancy, childbirth and perinatal period				
Tetanus neonatorum	A33	0-74		•
Obstetrical tetanus	A34	0-74		•
Pregnancy, childbirth and the puerperium	O00-O99	0-74	•	
Certain conditions originating in the perinatal period	P00-P96	0-74	•	
Congenital malformations				
Certain congenital malformations (neural tube defects)	Q00, Q01, Q05	0-74		•
Congenital malformations of the circulatory system (heart defects)	Q20-Q28	0-74	•	
Adverse effects of medical and surgical care				
Drugs, medicaments and biological substances causing adverse effects in therapeutic use	Y40-Y59	0-74	•	
Misadventures to patients during surgical and medical care	Y60-Y69, Y83-Y84	0-74	•	
Medical devices associated with adverse incidents in diagnostic and therapeutic use	Y70-Y82	0-74	•	
Injuries				
Transport Accidents	V01-V99	0-74		•
Accidental Injuries	W00-X39, X46-59	0-74		•
Intentional self-harm	X66-X84	0-74		•
Event of undetermined intent	Y16-Y34	0-74		•
Assault	X86-Y09	0-74		•
Alcohol related and drug-related deaths				
Alcohol specific disorders and poisonings	E24.4, F10, G31.2, G62.1, G72.1, I42.6, K29.2, K70, K85.2, K86.0, Q86.0, R78.0, X45, X65, Y15	0-74		•
Other alcohol related disorders	K73, K74.0-K74.2, K74.6	0-74		•
Drug disorders and poisonings	F11-F16, F18-F19, X40-X44, X85, Y10-Y14	0-74		•
Intentional self-poisoning by drugs	X60-X64	0-74		•
Provisional assignment of new diseases				
COVID-19	U07.1 - U07.2, U10.9	0-74		•

Table 8: Admissions – Alcohol Related Causes¹¹⁵

ICD-10 code	Description
E24.4	Alcohol induced Pseudo-Cushing's syndrome
E51.2	Wernicke's Encephalopathy
F10	Mental and Behavioural disorders due to use of alcohol
G31.2	Degeneration of the nervous system due to alcohol
G62.1	Alcoholic polyneuropathy
G72.1	Alcoholic myopathy
I42.6	Alcoholic cardiomyopathy
K29.2	Alcoholic gastritis
K70	Alcoholic liver disease
K85.2	Alcohol-induced acute pancreatitis
K86.0	Alcohol-induced chronic pancreatitis
O35.4	Maternal care for (suspected) damage to foetus from alcohol
P04.3	Foetus and newborn affected by maternal use of alcohol
Q86.0	Foetal alcohol syndrome (dysmorphic)
T51.0	Toxic effect of ethanol
T51.1	Toxic effect of methanol
T51.9	Toxic effect of alcohol, unspecified
X45	Accidental poisoning by and exposure to alcohol
X65	Intentional self-poisoning by and exposure to alcohol
Y15	Poisoning by and exposure to alcohol, undetermined intent
Y57.3	Alcohol deterrents
Y90	Evidence of alcohol involvement determined by blood alcohol level
Y91	Evidence of alcohol involvement determined by level intoxication
Z50.2	Alcohol rehabilitation
Z71.4	Alcohol abuse counselling and surveillance
Z72.1	Alcohol use

¹¹⁵ The definition for admissions due to alcohol related causes was updated for 2017/18 to include ICD Code K85.2: alcohol-induced chronic pancreatitis.

Table 9: Deaths – Alcohol Specific Causes

ICD-10 code	Description
E24.4	Alcohol-induced pseudo-Cushing's syndrome
F10	Mental and Behavioural disorders due to use of alcohol
G31.2	Degeneration of the nervous system due to alcohol
G62.1	Alcoholic polyneuropathy
G72.1	Alcohol myopathy
I42.6	Alcoholic cardiomyopathy
K29.2	Alcoholic gastritis
K70	Alcoholic liver disease
K85.2	Alcohol-induced acute pancreatitis
K86.0	Alcohol induced chronic pancreatitis
Q86.0	Fetal alcohol syndrome (dysmorphic)
R78.0	Excess alcohol blood levels
X45	Accidental poisoning by and exposure to alcohol
X65	Intentional self-poisoning by and exposure to alcohol
Y15	Poisoning by and exposure to alcohol, undetermined intent

Table 10: Admissions and Deaths – Drug Related Causes

ICD-10 code	Description
F11-16, F18-F19	Mental and Behavioural disorders due to drug use (excluding alcohol and tobacco)
X40-44	Accidental poisoning by drugs, medicaments and biological substances
X60-X64	Intentional self-poisoning by drugs, medicaments and biological substances
X85	Assault by drugs, medicaments and biological substances
Y10-14	Poisoning by drugs, medicaments and biological substances, undetermined intent

Table 11: Deaths – Drugs Misuse

ICD-10 code	Description
F11-16, F19	Mental and Behavioural disorders due to drug use (excluding alcohol, tobacco and volatile substances)
Deaths due to the following categories <i>and</i> where a drug controlled under the Misuse of Drugs Act 1971 was mentioned;	
X40-44	Accidental poisoning by drugs, medicaments and biological substances
X60-X64	Intentional self-poisoning by drugs, medicaments and biological substances
X85	Assault by drugs, medicaments and biological substances
Y10-14	Poisoning by drugs, medicaments and biological substances, undetermined intent

Table 12: Deaths – Smoking Attributable Causes ¹¹⁶

	ICD-10 Codes	Age	Attributable Proportion	
			Male	Female
Cancers which can be caused by smoking				
Upper respiratory sites	C10	35+	36%	31%
Upper respiratory sites	C11 & C30-C31	35+	23%	19%
Upper respiratory sites	C14	35+	59%	54%
Oesophagus	C15	35+	38%	31%
Stomach	C16	35+	16%	13%
Colorectal	C18-C20	35+	9%	7%
Liver	C22	35+	11%	9%
Pancreas	C25	35+	17%	14%
Larynx	C32	35+	69%	63%
Trachea, Lung, Bronchus	C33-C34	35+	73%	68%
Breast	C50	35+		3%
Cervical	C53	35+		15%
Kidney and Renal Pelvis	C64	35+	15%	12%
Kidney and Renal Pelvis	C65-C66	35+	35%	30%
Bladder	C67	35+	39%	34%
Myeloid leukaemia	C92	35+	12%	9%
Malignant leukaemia	C43-C44	50+	21%	
Cardiovascular diseases				
Ischaemic heart disease	I20-I25	35 - 54	38%	37%
	I20-I25	55 - 64	38%	39%
	I20-I25	65 - 74	16%	18%
	I20-I25	75+	15%	14%
Venous thromboembolism	I26, I80-I82	35+	7%	5%
Stroke	I60-I67	35+	11%	14%
Aortic aneurysm	I71	35+	20%	18%
Peripheral arterial disease	I73.9	35+	34%	29%
Respiratory diseases				
Tuberculosis	A15-A19	35+	9%	8%
Obstructive sleep apnoea	G47.3	35+	15%	13%
Influenza - microbiologically confirmed	J09, J10	35+	46%	42%
Influenza - clinically diagnosed	J11	35+	6%	5%
Pneumonia	J12-J18	35+	17%	15%
Chronic obstructive pulmonary disease	J40-J44, J47	35+	55%	48%
Adult asthma	J45-J46	35+	10%	8%
Idopathic Pulmonary fibrosis	J84.1	35+	9%	8%
Mental health				

¹¹⁶This definition is based on the latest Public Health England methodology ([https://www.gov.uk/government/statistics/local-tobacco-control-profiles-for-england-july-2021/local-tobacco-control-profiles-for-england-short-statistical-commentary-july-2021#:~:text=admissions%20\(new%20method\)-,Main%20findings,period%20\(2017%20to%202019\)\)](https://www.gov.uk/government/statistics/local-tobacco-control-profiles-for-england-july-2021/local-tobacco-control-profiles-for-england-short-statistical-commentary-july-2021#:~:text=admissions%20(new%20method)-,Main%20findings,period%20(2017%20to%202019)))) and estimates the impact of the latest NI smoking trends on observed mortality. The attributable proportions presented are representative of 2022 and will vary to a degree each year based on smoking prevalence statistics for the year in question.

Alzheimer's disease	G30	35+	8%	7%
Vascular dementia	F01	35+	5%	5%
All-cause dementia	F02, F03	35+	5%	5%
Depression	F32, F33	35+	10%	9%
Psychosis	F28, F29	35+	17%	15%
Schizophrenia	F20-F25	35+	18%	16%
Bulimia	F50.2	35+	19%	17%
Binge-eating disorder	F50.81	35+	28%	23%

Other diseases

Diabetes	E11	35+	10%	8%
Cataract	H25	35+	13%	10%
Age-related macular degeneration	H35.3-H52.4	35+	13%	12%
Hearing loss	H90, H91	35+	15%	13%
Barrett's oesophagus	K22.7	35+	17%	13%
Crohn's disease	K50	35+	12%	10%
Psoriasis	L40	35+	25%	20%
Rheumatoid arthritis	M05-M06	35+	15%	13%
Systemic lupus erythematosus	M32	35+	15%	12%
Low back pain	M54.5	35+	3%	2%
Chronic kidney disease	N18 (exc. N18.5)	35+	10%	8%
End-stage renal disease	N18.5	35+	23%	19%
Hip fracture	S72.0-S72.2	35+		5%
Surgical site infection	Y83, T81.4	35+	12%	11%

The table below lists the previous definition for all causes of death considered avoidable, with indication as to which are considered amenable, preventable or both. This list is for use with the additional indicators in [Appendix C](#).

Table 13: Amenable, Preventable and Avoidable Mortality (Previous ONS Definition).

Condition Group & Cause	ICD-10 Codes	Age	Amenable	Preventable
Infections				
Tuberculosis	A15-A19, B90	0-74	•	•
Selected invasive bacterial and protozoal infections	A38-A41, A46, A48.1, B50-B54, G00, G03, J02, L03	0-74	•	
Hepatitis C	B17.1, B18.2	0-74	•	•
Pertussis (whooping cough)	A37	0-14	•	•
Measles	B05	1-14	•	•
Rubella	B06	0-14		•
Other infections (Diphtheria, Tetanus, Poliomyelitis and Varicella)	A35, A36, A80, B01	0-19	•	•
Intestinal infections	A00-A09	0-14	•	

HIV/AIDS	B20-B24	All	•	•
Neoplasms				
Malignant neoplasm of lip, oral cavity and pharynx	C00-C14	0-74		•
Malignant neoplasm of oesophagus	C15	0-74		•
Malignant neoplasm of stomach	C16	0-74		•
Malignant neoplasm of colon and rectum	C18-C21	0-74	•	•
Malignant neoplasm of liver	C22	0-74		•
Malignant neoplasm of trachea, bronchus and lung	C33-C34	0-74		•
Malignant melanoma of skin	C43	0-74	•	•
Mesothelioma	C45	0-74		•
Malignant neoplasm of breast	C50	0-74	•	•
Malignant neoplasm of cervix uteri	C53	0-74	•	•
Malignant neoplasm of bladder	C67	0-74	•	
Malignant neoplasm of thyroid gland	C73	0-74	•	
Hodgkin's disease	C81	0-74	•	
Leukaemia	C91, C92.0	0-44	•	
Malignant neoplasm of testis	C62	0-74	•	
Malignant neoplasm of unspecified parts of uterus and body of uterus	C54-C55	0-44	•	
Benign neoplasms	D10-D36	0-74	•	
Nutritional, endocrine and metabolic				
Diabetes mellitus	E10-E14	0-74	•	•
Diseases of Thyroid	E00-E07	0-74	•	
Addison's Disease	E27.1	0-74	•	

Condition Group & Cause	ICD-10 Codes	Age	Amenable	Preventable
Drug use disorders				
Alcohol related diseases, excluding external causes	F10, G31.2, G62.1, I42.6, K29.2, K70, K73, K74 (excl. K74.3-K74.5), K86.0	0-74		•
Illicit drug use disorders	F11-F16, F18-F19	0-74		•
Neurological disorders				
Epilepsy and status epilepticus	G40-G41	0-74	•	
Cardiovascular diseases				
Rheumatic and other valvular heart disease	I01-I09	0-74	•	
Hypertensive diseases	I10-I15	0-74	•	
Ischaemic heart disease	I20-I25	0-74	•	•
DVT with pulmonary embolism	I26, I80.1-I80.3, I80.9, I82.9	0-74		•
Cerebrovascular diseases	I60-I69	0-74	•	
Aortic aneurysm and dissection	I71	0-74		•

Respiratory diseases					
Influenza (including swine flu)	J09-J11	0-74	•	•	
Pneumonia	J12-J18	0-74	•		
Chronic obstructive pulmonary disorder	J40-J44	0-74	•		•
Asthma	J45-J46	0-74	•		
Selected respiratory diseases	J00-J06, J20-J22, J30-J39	1-14	•		
Digestive disorders					
Gastric and duodenal ulcer	K25-K28	0-74	•		
Acute abdomen, appendicitis, intestinal obstruction, cholecystitis/lithiasis, pancreatitis, hernia	K35-K38, K40-K46, K80-K83, K85, K86.1-K86.9, K91.5	0-74	•		
Genitourinary disorders					
Nephritis and nephrosis	N00-N07, N17-N19, N25-N27	0-74	•		
Obstructive uropathy and prostatic hyperplasia	N13, N20-N21, N35, N40, N99.1	0-74	•		
Maternal and infant					
Complications of perinatal period	P00-P96, A33	All	•		
Congenital malformations of the circulatory system	Q20-Q28	0-74	•		
Spina Bifida	Q05	0-74			•
Unintentional injuries					
Transport Accidents	V01-V99	All			•
Accidental Injury	W00-X59	All			•
Intentional injuries					
Suicide and self inflicted injuries	X60-X84, Y10-Y34	All			•
Homicide/Assault	X85-Y09, U50.9	All			•
Misadventures to patients during surgical and medical care	Y60-Y69, Y83-Y84	All	•		•

Also available for the Health & Social Care Inequalities Monitoring System (HSCIMS)

Life Expectancy in Northern Ireland

Presenting the latest official estimates of life expectancy in Northern Ireland and an examination of the causes that contribute to the change in life expectancy over time, as well as the differentials between gender and deprivation classification and across HSC Trusts and Local Government Districts. The latest figures for life expectancy at 65, healthy life expectancy and disability-free life expectancy are also included.

www.health-ni.gov.uk/articles/life-expectancy-northern-ireland

Making Life Better

Monitoring the Wider Social Determinants of Health & Wellbeing Key Indicators

Monitoring of the key indicators of the wider social determinant of health & wellbeing set out against each of the themes contained in the making life better strategic framework (Annual).

www.health-ni.gov.uk/articles/social-determinants-health-statistics

Produced by:

Public Health Information & Research Branch, Information Analysis Directorate

Department of Health

Castle Buildings, Belfast BT4 3SQ

Tel: 02890522501

Email: healthinequalities@health-ni.gov.uk

www.health-ni.gov.uk/articles/health-inequalities-statistics